

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : <b>C07D 211/66, A61K 31/445, C07D 309/08, A61K 31/35, C07D 401/06, 409/06, 401/12, 413/04, 407/06, A61P 19/00, 19/02</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 00/44723</b></p> <p>(43) International Publication Date: <b>3 August 2000 (03.08.00)</b></p>
<p>(21) International Application Number: <b>PCT/US00/01864</b></p> <p>(22) International Filing Date: <b>27 January 2000 (27.01.00)</b></p> <p>(30) Priority Data: <b>09/238,038      27 January 1999 (27.01.99)      US</b></p> <p>(71) Applicant: <b>AMERICAN CYANAMID COMPANY [US/US];</b> <b>Five Giralda Farms, Madison, NJ 07940-0874 (US).</b></p> <p>(72) Inventors: <b>LEVIN, Jeremy, Ian; 19 Long Meadow Drive, New City, NY 10956 (US). VENKATESAN, Aranapakam, Mudumbai; 97-07, 63rd Road #9K, Rego Park, NY 11374 (US). CHEN, James, Ming; 7 Sergeant David Stoddard Court, Bedminster, NJ 07921 (US). ZASK, Arie; 21 East 90 Street, New York, NY 10128 (US). SANDANAYAKA, Vincent, Premarana; #1 Earl Court, Wesley Hills, NY 10952 (US). DU, Mila, Ti; 17 Heather Hill Lane, Suffern, NY 10901 (US). BAKER, Jannie, Lea; 127 Rockinchair Road, White Plains, NY 10607 (US).</b></p> <p>(74) Agents: <b>BARRETT, Rebecca, R.; American Home Products Corporation, Patent Law Dept. - 2B, One Campus Drive, Parsippany, NJ 07054 (US) et al.</b></p>	<p>(81) Designated States: <b>AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</b></p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	

(54) Title: **ALKYNYL CONTAINING HYDROXAMIC ACID DERIVATIVES, THEIR PREPARATION AND THEIR USE AS MATRIX METALLOPROTEINASE (MMP) INHIBITORS/TNF-ALPHA CONVERTING ENZYME (TACE) INHIBITORS**

(57) Abstract

Compounds of formula (I) wherein R<sub>1</sub> is hydrogen, aryl, heteroaryl, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, or -C<sub>4</sub>-C<sub>8</sub>-cycloheteroalkyl; R<sub>2</sub> and R<sub>3</sub> are hydrogen, alkyl of 1-6 carbon atoms, -CN, or CCH;



R<sub>8</sub>, R<sub>9</sub>, R<sub>10</sub> and R<sub>11</sub> are hydrogen, aryl or heteroaryl, cycloalkyl of 3-6 carbon atoms, -C<sub>4</sub>-C<sub>8</sub>-cycloheteroalkyl, alkyl of 1-18 carbon atoms, alkenyl of 2-18 carbon atoms, alkynyl of 2-18 carbon atoms; with the proviso that one of the pairs R<sub>8</sub> and R<sub>9</sub>, R<sub>9</sub> and R<sub>10</sub> or R<sub>10</sub> and R<sub>11</sub>, together with the carbon atom or atoms to which they are attached, form a cycloalkyl ring of 3-6 carbon atoms, or a -C<sub>4</sub>-C<sub>8</sub>-cycloheteroalkyl ring; R<sub>12</sub> is hydrogen, aryl or heteroaryl, cycloalkyl of 3-6 carbon atoms; -C<sub>4</sub>-C<sub>8</sub>-cycloheteroalkyl, or alkyl of 1-6 carbon atoms; A, X are O, S, SO, SO<sub>2</sub>, NR<sub>7</sub>, or CH<sub>2</sub>; Y is aryl or heteroaryl, with the proviso that A and X are not bonded to adjacent atoms of Y; and n is 0-2; or a pharmaceutically acceptable salt thereof; useful in the treatment of arthritis, tumor metastasis, tissue ulceration, abnormal wound healing, periodontal disease, bone disease, diabetes (insulin resistance) and HIV infection.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

ALKYNYL CONTAINING HYDROXAMIC ACID DERIVATIVES, THEIR PREPARATION AND THEIR  
USE AS MATRIX METALLOPROTEINASE (MMP) INHIBITORS / TNF-ALPHA CONVERTING  
ENZYME (TACE) INHIBITORS

**FIELD OF INVENTION**

5        This invention relates to acetylenic hydroxamic acids which act as inhibitors  
of TNF- $\alpha$  converting enzyme (TACE). The compounds of the present invention are  
useful in disease conditions mediated by TNF- $\alpha$ , such as rheumatoid arthritis,  
osteoarthritis, sepsis, AIDS, ulcerative colitis, multiple sclerosis, Crohn's disease and  
degenerative cartilage loss.

10

**BACKGROUND OF THE INVENTION**

Matrix metalloproteinases (MMPs) are a group of enzymes that have been  
implicated in the pathological destruction of connective tissue and basement  
membranes. These zinc containing endopeptidases consist of several subsets of  
15 enzymes including collagenases, stromelysins and gelatinases. Of these classes, the  
gelatinases have been shown to be the MMPs most intimately involved with the  
growth and spread of tumors. It is known that the level of expression of gelatinase is  
elevated in malignancies, and that gelatinase can degrade the basement membrane  
which leads to tumor metastasis. Angiogenesis, required for the growth of solid  
20 tumors, has also recently been shown to have a gelatinase component to its pathology.  
Furthermore, there is evidence to suggest that gelatinase is involved in plaque rupture  
associated with atherosclerosis. Other conditions mediated by MMPs are restenosis,  
MMP-mediated osteopenias, inflammatory diseases of the central nervous system,  
skin aging, tumor growth, osteoarthritis, rheumatoid arthritis, septic arthritis, corneal  
25 ulceration, abnormal wound healing, bone disease, proteinuria, aneurysmal aortic  
disease, degenerative cartilage loss following traumatic joint injury, demyelinating  
diseases of the nervous system, cirrhosis of the liver, glomerular disease of the  
kidney, premature rupture of fetal membranes, inflammatory bowel disease,  
periodontal disease, age related macular degeneration, diabetic retinopathy,  
30 proliferative vitreoretinopathy, retinopathy of prematurity, ocular inflammation,  
keratoconus, Sjogren's syndrome, myopia, ocular tumors, ocular angiogenesis/neo-


vascularization and corneal graft rejection. For recent reviews, see: (1) Recent Advances in Matrix Metalloproteinase Inhibitor Research, R. P. Beckett, A. H. Davidson, A. H. Drummond, P. Huxley and M. Whittaker, Research Focus, Vol. 1, 16-26, (1996), (2) Curr. Opin. Ther. Patents (1994) 4(1): 7-16, (3) Curr. Medicinal Chem. (1995) 2: 743-762, (4) Exp. Opin. Ther. Patents (1995) 5(2): 1087-110, (5) 5 Exp. Opin. Ther. Patents (1995) 5(12): 1287-1196: (6) Exp. Opin. Ther. Patents (1998) 8(3): 281-259.

TNF- $\alpha$  converting enzyme (TACE) catalyzes the formation of TNF- $\alpha$  from membrane bound TNF- $\alpha$  precursor protein. TNF- $\alpha$  is a pro-inflammatory cytokine that is believed to have a role in rheumatoid arthritis [Shire, M. G.; Muller, G. W. *Exp. Opin. Ther. Patents* **1998**, 8(5), 531; Grossman, J. M.; Brahn, E. J. *Women's Health* **1997**, 6(6), 627; Isomaki, P.; Punnonen, J. *Ann. Med.* **1997**, 29, 499; Camussi, G.; Lupia, E. *Drugs*, **1998**, 55(5), 613.] septic shock [Mathison, et. al. *J. Clin. Invest.* **1988**, 81, 1925; Miethke, et. al. *J. Exp. Med.* **1992**, 175, 91.], graft rejection [Piguet, P. F.; Grau, G. E.; et. al. *J. Exp. Med.* **1987**, 166, 1280.], cachexia [Beutler, B.; Cerami, A. *Ann. Rev. Biochem.* **1988**, 57, 505.], anorexia, inflammation [Ksontini, R.; MacKay, S. L. D.; Moldawer, L. L. *Arch. Surg.* **1998**, 133, 558.], congestive heart failure [Packer, M. *Circulation*, **1995**, 92(6), 1379; Ferrari, R.; Bachetti, T.; et. al. *Circulation*, **1995**, 92(6), 1479.], post-ischaemic reperfusion injury, inflammatory disease of the central nervous system, inflammatory bowel disease, insulin resistance [Hotamisligil, G. S.; Shargill, N. S.; Spiegelman, B. M.; et. al. *Science*, **1993**, 259, 87.] and HIV infection [Peterson, P. K.; Gekker, G.; et. al. *J. Clin. Invest.* **1992**, 89, 574; Pallares-Trujillo, J.; Lopez-Soriano, F. J. Argiles, J. M. *Med. Res. Reviews*, **1995**, 15(6), 533.], in addition to its well-documented antitumor properties [Old, L. *Science*, **1985**, 230, 630.]. For example, research with anti-TNF- $\alpha$  antibodies and transgenic animals has demonstrated that blocking the formation of TNF- $\alpha$  inhibits the progression of arthritis [Rankin, E.C.; Choy, E.H.; Kassimos, D.; Kingsley, G.H.; Sopwith, A.M.; Isenberg, D.A.; Panayi, G.S. *Br. J. Rheumatol.* **1995**, 34, 334; *Pharmaprojects*, **1996**, Therapeutic Updates 17 (Oct.), au197-M2Z.]. This observation has recently been extended to humans as well as described in "TNF- $\alpha$  in Human Diseases", *Current Pharmaceutical Design*, **1996**, 2, 662.


-3-

It is expected that small molecule inhibitors of TACE would have the potential for treating a variety of disease states. Although a variety of TACE inhibitors are known, many of these molecules are peptidic and peptide-like which suffer from bioavailability and pharmacokinetic problems. In addition, many of these molecules are non-selective, being potent inhibitors of matrix metalloproteinases and, in particular, MMP-1. Inhibition of MMP-1 (collagenase 1) has been postulated to cause joint pain in clinical trials of MMP inhibitors [*Scrip*, 1998, 2349, 20]. Long acting, selective, orally bioavailable non-peptide inhibitors of TACE would thus be highly desirable for the treatment of the disease states discussed above.

Sulfone hydroxamic acid inhibitors of MMPs, of general structure I have been disclosed [Burgess, L.E.; Rizzi, J.P.; Rawson, D.J. **Eur Patent Appl. 818442**. Groneberg, R.D.; Neuenschwander, K.W.; Djuric, S.W.; McGeehan, G.M.; Burns, C.J.; Condon, S.M.; Morrisette, M.M.; Salvino, J.M.; Scotese, A.C.; Ullrich, J.W. **PCT Int. Appl. WO 97/24117**. Bender, S.L.; Broka, C.A.; Campbell, J.A.; Castelhana, A.L.; Fisher, L.E.; Hendricks, R.T.; Sarma, K. **Eur. Patent Appl. 780386**. Venkatesan, A. M.; Grosu, G. T.; Davis, J. M.; Hu, B.; O'Dell, M. J. **PCT Int. Appl. WO 98/38163**.]. An exemplification of this class of MMP inhibitor is RS-130830, shown below.



I




RS-130830

20


Within the sulfone-hydroxamic acid class of MMP inhibitor, the linker between the sulfone and hydroxamic acid moieties has been extended to three carbons (I, n = 2) without significant loss in potency [Barta, T. E.; Becker, D. P.; Villamil, C. I.;

Freskos, J. N.; Mischke, B. V.; Mullins, P. B.; Heintz, R. M.; Getman, D. P.; McDonald, J. J. **PCT Int. Appl. WO 98/39316**. McDonald, J. J.; Barta, T. E.; Becker, D. P.; Bedell, L. J.; Rao, S. N.; Freskos, J. N.; Mischke, B. V. **PCT Int. Appl. WO 98/38859**.]


- 5 Piperidine sulfone hydroxamic acids, **II** ( $n = 1$ ) have been reported [Becker, D. P.; Villamil, C. I.; Boehm, T. L.; Getman, D. P.; McDonald, J. J.; DeCrescenzo, G. A. **PCT Int. Appl. WO 98/39315**.]. Similar piperidine derivatives in which the methylene linking the piperidine ring to the sulfone has been deleted (**II**,  $n = 0$ ) have been reported [Venkatesan, A. M.; Grosu, G. T.; Davis, J. M.; Baker, J. L. **PCT Int. Appl. WO 98/37877**.].



- Sulfone-hydroxamic acids **III**, in which a hydroxyl group has been placed alpha to the hydroxamic acid, have been disclosed [Freskos, J. N.; Boehm, T. L.; Mischke, B. V.; Heintz, R. M.; McDonald, J. J.; DeCrescenzo, G. A.; Howard, S. C. **PCT Int. Appl. WO 98/39326**. Robinson, R. P. **PCT Int. Appl. WO 98/34915**.].




- 20 Sulfone-based MMP inhibitors of general structure **IV**, which utilize a thiol as the zinc chelator, have been reported [Freskos, J.N.; Abbas, Z.S.; DeCrescenzo, G.A.; Getman, D.P.; Heintz, R.M.; Mischke, B.V.; McDonald, J.J. **PCT Int. Appl. WO 98/03164**.].




-5-

Inhibitors of stromelysin with general structure V have been disclosed [Shuker, S.B.; Hajduk, P.J.; Meadows, R.P.; Fesik, S.W. *Science*, **1996**, 274, 1531-1534. Hajduk, P.J.; Sheppard, G.; Nettesheim, D.G.; Olejniczak, E.T.; Shuker, S.B.; Meadows, R.P.; Steinman, D.H.; Carrera, Jr., G.M.; Marcotte, P.A.; Severin, J.; Walter, K.; Smith, H.; Gubbins, E.; Simmer, R.; Holzman, T.F.; Morgan, D.W.; Davidsen, S.K.; Summers, J.B.; Fesik, S.W. *J. Am. Chem. Soc.* **1997**, 119, 5818-5827. Olejniczak, E.T.; Hajduk, P.J.; Marcotte, P.A.; Nettesheim, D.G.; Meadows, R.P.; Edalji, R.; Holzman, T.F.; Fesik, S.W. *J. Am. Chem. Soc.* **1997**, 119, 5828-5832. Fesik, S. W.; Summers, J. B.; Davidsen, S. K.; Sheppard, G. S.; Steinman, D. H.; Carrera, G. M.; Florjancic, A.; Holms, J. H. *PCT Int. Appl.* **WO 97/18188**.]



Salah et al., Liebigs Ann. Chem. 195, (1973) discloses some aryl substituted thio and aryl substituted sulfonyl acetohydroxamic acid derivatives of general formula 1. These compounds were prepared to study the Mannich reaction. Subsequently, they were tested for their fungicidal activity.




Some sulfone carboxylic acids are disclosed in U.S. patent 4,933,367. Those compounds were shown to exhibit hypoglycemic activity.

-6-

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to novel, low molecular weight, non-peptide inhibitors of matrix metalloproteinases (MMPs) and TNF- $\alpha$  converting enzyme (TACE) for the treatment of arthritis, tumor metastasis, tissue ulceration, abnormal wound healing, periodontal disease, bone disease, diabetes (insulin resistance) and HIV infection.

In accordance with this invention there is provided a group of compounds of general formula I:



wherein:

$R_1$  is hydrogen, aryl, heteroaryl, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, or -C4-C8-cycloheteroalkyl;

15

$R_2$  and  $R_3$  are each, independently, hydrogen, alkyl of 1-6 carbon atoms, -CN, or -CCH<sub>3</sub>;

$R_4$  is hydrogen, aryl, aralkyl, heteroaryl, heteroaralkyl, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 1-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, -C(O)- $R_1$ , -SO<sub>2</sub>- $R_1$ , -C(O)-NHR<sub>1</sub>, -C(O)NR<sub>5</sub>R<sub>6</sub>, -C(O)R<sub>1</sub>NR<sub>5</sub>R<sub>6</sub>, -C(O)-OR<sub>1</sub>, -C(NH)-NH<sub>2</sub>.

20

$R_5$  and  $R_6$  are each, independently, hydrogen, alkyl of 1-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, aryl, aralkyl, heteroaryl, heteroaralkyl or -C4-C8-cycloheteroalkyl;

25



-7-

$R_8$ ,  $R_9$ ,  $R_{10}$ , and  $R_{11}$  are each, independently, hydrogen, aryl or heteroaryl, cycloalkyl of 3-6 carbon atoms, -C4-C8-cycloheteroalkyl, alkyl of 1-18 carbon atoms, alkenyl of 2-18 carbon atoms, alkynyl of 2-18 carbon atoms; with the proviso that one of the pairs  $R_8$  and  $R_9$ ,  $R_9$  and  $R_{10}$  or  $R_{10}$  and  $R_{11}$ , together with the carbon atom or atoms to which they are attached, form a cycloalkyl ring of 3-6 carbon atoms, or a -C4-C8-cycloheteroalkyl ring;

$R_{12}$  is hydrogen, aryl or heteroaryl, cycloalkyl of 3-6 carbon atoms, -C4-C8-cycloheteroalkyl, or alkyl of 1-6 carbon atoms;

10 A is O, S, SO,  $SO_2$ ,  $NR_7$ , or  $CH_2$ ;

X is O, S, SO,  $SO_2$ ,  $NR_7$ , or  $CH_2$ ;

Y is aryl or heteroaryl, with the proviso that A and X are not bonded to adjacent atoms of Y; and

n is 0-2; or a pharmaceutically acceptable salt thereof.

In some preferred aspects of the invention, Y is phenyl, pyridyl, thienyl, furanyl, imidazolyl, triazolyl or thiadiazolyl, with the proviso that A and X are not bonded to adjacent atoms of Y.

In still other preferred embodiments of the invention Y is phenyl, thienyl or furanyl.

25 In accordance with certain preferred embodiments of the invention  $R_8$  and  $R_9$ , together with the carbon atom to which they are attached form a  $C_4$ - $C_8$  cycloheteroalkyl ring and K is  $NR_7$ .

The most preferred matrix metalloproteinase and TACE inhibiting compounds of this invention are:

1-(4-Bromo-benzyl)-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide;

4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid hydroxyamide;

- 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-chloro-benzyl)-piperdine-4-carboxylic acid hydroxyamide;
- 1-Benzyl-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperdine-4-carboxylic acid hydroxamide;
- 5 1-(4-Bromo-benzyl)-4-(4-pent-2-ynyloxy-benzenesulfonyl)-piperdine-4-carboxylic acid hydroxyamide;
- 1-(4-Bromo-benzyl)-4-(4-oct-2-ynyloxy-benzenesulfonyl)-piperdine-4-carboxylic acid hydroxyamide;
- 10 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-fluoro-benzyl)-piperdine-4-carboxylic acid hydroxyamide;
- 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-cyano-benzyl)-piperidine-4-carboxylic acid hydroxamide;
- 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-methyl-benzyl)-piperidine-4-carboxylic acid hydroxamide;
- 15 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(3,4- dichloro-benzyl)-piperidine-4-carboxylic acid hydroxamide;
- 1-(4-Bromo-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperdine-4-carboxylic acid hydroxyamide;
- 20 1-(4-Bromo-benzyl)-4-[4-(4-piperdin-4-yl-but-2-ynyloxy)-benzenesulfonyl]-piperdine-4-carboxylic acid hydroxyamide;
- 1-(4-Bromo-benzyl)-4-[4-(4-morpholin-4-yl-but-2-ynyloxy)-benzene-sulfonyl]-piperdine-4-carboxylic acid hydroxyamide;
- 4-(4-But-2-ynyloxy-phenylsulfanyl)-4-hydroxycarbamoyl-piperidine-1-carboxylic acid tert-butyl ester;
- 25 4-(4-But-2-ynyloxy-phenylsulfanyl)-piperidine-4-carboxylic acid hydroxyamide
- 1-(4-Bromo-benzyl)-4-(4-but-2-ynyloxy-phenylsulfanyl)-piperidine-4-carboxylic acid hydroxyamide;
- 30 4-(4-But-2-ynyloxy-phenylsulfanylmethyl)-tetrahydro-pyran-4-carboxylic acid hydroxyamide;

- 4-(4-But-2-ynyloxy-benzenesulfonylmethyl)-tetrahydro-pyran-4-carboxylic acid hydroxyamide;
- 4-(4-But-2-ynyloxy-benzenesulfinylmethyl)-tetrahydro-pyran-4-carboxylic acid hydroxyamide;
- 5 4-{[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxytetrahydro-2H-pyran-4-carboxamide;
- 1-benzyl-4-{[3-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxy-4-piperidine carboxamide;
- 10 4-{[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxy-1-isopropyl-4-piperidine carboxamide;
- 4-{[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxy-1-(3-pyridinylmethyl)-4-piperidine carboxamide;
- 3-{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-ethyl-N-hydroxy-3-piperidine-carboxamide;
- 15 3-{[4-(2-butynyloxy)phenyl]sulfonyl}-1-(4-chlorobenzyl)-N-hydroxy-3-piperidinecarboxamide;
- 4-{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid hydroxyamide;
- 4-{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-(3-pentanyl)-piperidine-4-carboxylic acid hydroxyamide;
- 20 1-(4-Methoxy-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide;
- 1-(4-Chloro-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide;
- 25 tert-butyl-4-({[4-(2-butynyloxy)phenyl]sulfanyl}methyl)-4-[(hydroxyamino)-carbonyl]-1-piperidinecarboxylate;
- 4-({[4-(But-2-ynyloxy)phenyl]thio}methyl)-N-hydroxypiperidine-4-carboxamide;
- tert-Butyl-4-({[4-(2-butynyloxy)phenyl]sulfinyl}methyl)-4-[(hydroxyamino)-carbonyl]-1-piperidinecarboxylate;
- 30

- 4-[[[4-(2-Butynyloxy)phenyl]sulfinyl]methyl]-N-hydroxy-4-piperidine-carboxamide;
- tert-Butyl-4-({[4-(but-2-ynyloxy)phenyl]sulfonyl}methyl)-4-[(hydroxyamino)-carbonyl]piperidine-1-carboxylate;
- 5 tert-butyl-4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-4-[(hydroxyamino)-carbonyl]-1-piperidinecarboxylate;
- 1-Acetyl-4-[[[4-(2-butynyloxy)phenyl]sulfonyl]methyl]-N-hydroxy-4-piperidinecarboxamide;
- 1-(2-Butynyl)-4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-4-piperidinecarboxamide hydrochloride;
- 10 N-1-(tert-Butyl)-4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-N-4-hydroxy-1,4-[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-N-4-hydroxy-1,4-l[sulfonyl}-methyl)-N-4-hydroxy-1,4-piperidinedicarboxamide;
- Methyl 4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-4-[(hydroxyamino)-carbonyl]-1-piperidinecarboxylate;
- 15 Benzyl 4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-4-[(hydroxyamino)-carbonyl]-1-piperidinecarboxylate;
- 1-Benzyl-4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-4-butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-4-piperidinecarboxamide;
- 20 4-({[4-(2-Butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxamide;
- 4-({[4-(2-Butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-1-[3-hydroxy-2-(hydroxymethyl)-2-methylpropanoyl]-4-piperidinecarboxamide;
- 1-[Amino(imino)methyl]-4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-4-l]-4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-4-oxy)phenyl]sulfonyl}methyl)-N-hydroxy-4-piperidinecarboxamide;
- 25 4-({[4-(2-Butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-1-(4-hydroxy-2-butynyl)-henyl]sulfonyl}methyl)-N-hydroxy-1-(4-hydroxy-2-butynyl)-4-piperidinecarboxamide;

- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-1-ethyl-N-hydroxypiperidine-4-carboxamide trifluoroacetic acid salt;
- 2-chloro-5-(chloromethyl) thiophene4-([4-(But-2-ynyloxy)phenyl]-sulfonyl)-methyl)-1-[(5-chlorothien-2-yl)methyl]-N-hydroxypiperidine-4-carboxamide trifluoroacetic acid salt;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(pyridin-4-ylmethyl)piperidine-4-carboxamide trifluoroacetic acid salt;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(pyridin-3-ylcarbonyl)piperidine-4-carboxamide trifluoroacetic acid salt;
- 1-Benzoyl-4-([4-(but-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxypiperidine-4-carboxamide;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(thien-2-ylcarbonyl)piperidine-4-carboxamide;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-1-ethyl-N-4-hydroxypiperidine-1,4-dicarboxamide;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-4-hydroxy-N-1-phenylpiperidine-1,4-dicarboxamide;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-1-,N-1-diethyl-N-4-hydroxypiperidine-1,4-dicarboxamide;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(morpholin-4-ylcarbonyl)piperidine-4-carboxamide;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-4-hydroxy-N-1-methyl-N-1-phenylpiperidine-1,4-dicarboxamide;
- Octyl-4-([4-(but-2-ynyloxy)phenyl]sulfonyl)methyl)-4-[(hydroxyamino)-carbonyl]piperidine-1-carboxylate;
- 4-Methoxyphenyl4-([4-(but-2-ynyloxy)phenyl]sulfonyl)methyl)-4-[(hydroxyamino) carbonyl]piperidine-1-carboxylate;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(phenylsulfonyl)piperidine-4-carboxamide;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-[(1-methyl-1H-imidazol-4-yl)sulfonyl]piperidine-4-carboxamide;


-12-

- 1-[2-(Benzylamino)acetyl]-4-({[4-(but-2-ynyloxy)phenyl]-sulfonyl}methyl)-N-hydroxypiperidine-4-carboxamide;
- 4-({[4-(But-2-ynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-1-(2-morpholin-4-ylacetyl)piperidine-4-carboxamide;
- 5 4-({[4-(But-2-ynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-1-[2-(4-methylpiperazin-1-yl)acetyl]piperidine-4-carboxamide;
- 1-Acetyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid hydroxamide;
- 10 1-Benzoyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid hydroxamide;
- 1-(4-Methoxybenzoyl)-4-(4-but-2-ynyloxy benzenesulfonyl)piperidine-4-carboxylic acid hydroxamide;
- 4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(pyrrolidine-1-carbonyl)-4-piperidinecarboxamide;
- 15 Ethyl 4-(4-but-2-ynyloxybenzenesulfonyl)-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxylate;
- 4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(trifluoromethyl)sulfonyl]-4-piperidinecarboxamide;
- 4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(3-pyridinylcarbonyl)-4-piperidinecarboxamide;
- 20 4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(2-thienylcarbonyl)-4-piperidinecarboxamide;
- 4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(4-methoxyphenyl)sulfonyl]-4-piperidinecarboxamide;
- 25 4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxamide;
- Tert-butyl-4-({[4-(2-butynyloxy)phenyl]sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxalate;
- 4-({[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxy-4-piperidinecarboxamide hydrochloride;
- 30

-13-

- Methyl ({4-{{4-(2-butynyloxy)phenyl}sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinyl}methyl)benzoate hydrochloride;  
 4-({4-{{4-(2-butynyloxy)phenyl}sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinyl}methyl)benzoic acid hydrochloride;  
 5 1-[4-(Aminocarbonyl)benzyl]-4-{{4-(2-butynyloxy)phenyl}sulfonyl}-N-hydroxy-4-piperidinecarboxamide hydrochloride;  
 Tert-butyl 4-{{4-(but-2-ynyloxy)phenyl}sulfinyl}-4-[(hydroxyamino)-carbonyl]piperidine-1-carboxalate;  
 4-(4-(But-2-ynyloxy-benzenesulfinyl)-piperidine-4-carboxylic acid  
 10 hydroxamide hydrochloride; and  
 1-(4-Bromo-benzyl)-4-(4-But-2-ynyloxy-benzenesulfinyl)-piperidine-4-carboxylic acid hydroxamide hydrochloride;  
 and pharmaceutical salts thereof.

- 15 Heteroaryl, as used throughout, is a 5-10 membered mono- or bicyclic aromatic ring having from 1-3 heteroatoms selected from N, NR<sub>7</sub>, S and O. Heteroaryl is preferably




-14-

wherein K is defined as O, S or -NR<sub>7</sub>, and R<sub>7</sub> is hydrogen, aryl, aralkyl, heteroaryl, heteroaralkyl, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 1-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, -C(O)-R<sub>1</sub>, -SO<sub>2</sub>-R<sub>1</sub>, -C(O)-NHR<sub>1</sub>, -C(O)NR<sub>5</sub>R<sub>6</sub>, -C(O)R<sub>1</sub>, NR<sub>5</sub>R<sub>6</sub>, -C(O)-OR<sub>1</sub>, -C(NH)-NH<sub>2</sub>.

Preferred heteroaryl rings include pyrrole, furan, thiophene, pyridine, pyrimidine, pyridazine, pyrazine, triazole, pyrazole, imidazole, isothiazole, thiazole, isoxazole, oxazole, indole, isoindole, benzofuran, benzothiophene, quinoline, isoquinoline, quinoxaline, quinazoline, benzotriazole, indazole, benzimidazole, benzothiazole, benzisoxazole, and benzoxazole. Heteroaryl groups of the present invention may be mono or disubstituted.

-C<sub>4</sub>-C<sub>8</sub>-cycloheteroalkyl is defined as



wherein K is O, S or NR<sub>7</sub> and R<sub>7</sub> is as defined before. Preferred heterocycloalkyl rings include piperidine, piperazine, morpholine, tetrahydropyran, tetrahydrofuran or pyrrolidine. Heterocycloalkyl groups of the present invention may optionally be mono- or disubstituted.



-15-

Aryl, as used herein refers to phenyl or naphthyl aromatic rings which may, optionally be mono- or di- substituted.

Alkyl, alkenyl, alkynyl, and perfluoroalkyl include both straight chain as well  
 5 as branched moieties. Alkyl, alkenyl, alkynyl, and cycloalkyl groups may be unsubstituted (carbons bonded to hydrogen, or other carbons in the chain or ring) or may be mono- or poly-substituted.

Aralkyl as used herein refers to a substituted alkyl group, -alkyl-aryl, wherein alkyl is lower alkyl and preferably from 1-3 carbon atoms, and aryl is as previously  
 10 defined.

Heteroaralkyl as used herein refers to a substituted alkyl group, alkyl-heteroaryl wherein alkyl is lower alkyl and preferably from 1-3 carbon atoms, and heteroaryl is as previously defined.

15 Halogen means bromine, chlorine, fluorine, and iodine.

Suitable substituents of aryl, aralkyl, heteroaryl, heteroaralkyl, alkyl, alkenyl, alkynyl and cycloalkyl include, but are not limited to halogen, alkyl of 1-6 carbon atoms; alkenyl of 2-6 carbon atoms; alkynyl of 2-6 carbon atoms, cycloalkyl  
 20 of 3-6 carbon atoms, -OR<sub>5</sub>, =O, -CN, -COR<sub>5</sub>, perfluoroalkyl of 1-4 carbon atoms, -O-perfluoroalkyl of 1-4 carbon atoms, -CONR<sub>5</sub>R<sub>6</sub>, -S(O)<sub>n</sub>R<sub>5</sub>, -OPO(OR<sub>5</sub>)OR<sub>6</sub>, -PO(OR<sub>5</sub>)R<sub>6</sub>, -OC(O)OR<sub>5</sub>, -OR<sub>5</sub>NR<sub>5</sub>R<sub>6</sub>, -OC(O)NR<sub>5</sub>R<sub>6</sub>, -C(O)NR<sub>5</sub>OR<sub>6</sub>, -COOR<sub>5</sub>, -SO<sub>3</sub>H, -NR<sub>5</sub>R<sub>6</sub>, -N[(CH<sub>2</sub>)<sub>2</sub>]<sub>2</sub>NR<sub>5</sub>, -NR<sub>5</sub>COR<sub>6</sub>, -NR<sub>5</sub>COOR<sub>6</sub>, -SO<sub>2</sub>NR<sub>5</sub>R<sub>6</sub>, -NO<sub>2</sub>, -N(R<sub>5</sub>)SO<sub>2</sub>R<sub>6</sub>, -NR<sub>5</sub>CONR<sub>5</sub>R<sub>6</sub>,  
 25 -NR<sub>5</sub>C(=NR<sub>6</sub>)NR<sub>5</sub>R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)N(SO<sub>2</sub>R<sub>5</sub>)R<sub>6</sub>, -NR<sub>5</sub>C(=NR<sub>6</sub>)N(C=OR<sub>5</sub>)R<sub>6</sub>, -tetrazol-5-yl, -SO<sub>2</sub>NHCN, -SO<sub>2</sub>NHCONR<sub>5</sub>R<sub>6</sub>, phenyl, heteroaryl or -C<sub>4</sub>-C<sub>8</sub>-cycloheteroalkyl;

wherein -NR<sub>5</sub>R<sub>6</sub> may form a pyrrolidine, piperidine, morpholine, thiomorpholine, oxazolidine, thiazolidine, pyrazolidine, piperazine, or azetidine ring;

30

-16-

$R_5$  and  $R_6$  are each, independently, hydrogen, alkyl of 1-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, aryl, aralkyl, heteroaryl, heteroaralkyl or -C<sub>4</sub>-C<sub>8</sub>-cycloheteroalkyl;

- 5  $R_7$  is hydrogen, aryl, heteroaryl, alkyl of 1-6 carbon atoms or cycloalkyl of 3-6 carbon atoms, -C(O)- $R_1$ , -SO<sub>2</sub>- $R_1$ , -C(O)-NHR<sub>1</sub>, -C(O)-OR<sub>1</sub>, -C(NH)-NH<sub>2</sub>;

and n is 0-2.

- 10 When a moiety contains more than one substituent with the same designation each of those substituents may be the same or different.

Pharmaceutically acceptable salts can be formed from organic and inorganic acids, for example, acetic, propionic, lactic, citric, tartaric, succinic, fumaric, maleic, 15 malonic, mandelic, malic, phthalic, hydrochloric, hydrobromic, phosphoric, nitric, sulfuric, methanesulfonic, naphthalenesulfonic, benzenesulfonic, toluenesulfonic, camphorsulfonic, and similarly known acceptable acids when a compound of this invention contains a basic moiety. Salts may also be formed from organic and inorganic bases, preferably alkali metal salts, for example, sodium, lithium, or 20 potassium, when a compound of this invention contains an acidic moiety.


The compounds of this invention may contain an asymmetric carbon atom and some of the compounds of this invention may contain one or more asymmetric centers and may thus give rise to optical isomers and diastereomers. While shown without respect to stereochemistry, the present invention includes such optical 25 isomers and diastereomers; as well as the racemic and resolved, enantiomerically pure R and S stereoisomers; as well as other mixtures of the R and S stereoisomers and pharmaceutically acceptable salts thereof. It is recognized that one optical isomer, including diastereomer and enantiomer, or stereoisomer may have favorable properties over the other. Thus when disclosing and claiming the invention, when one 30 racemic mixture is disclosed, it is clearly contemplated that both optical isomers, including diastereomers and enantiomers, or stereoisomers substantially free of the other are disclosed and claimed as well.

-17-

The compounds of this invention are shown to inhibit the enzymes MMP-1, MMP-9, MMP-13 and TNF- $\alpha$  converting enzyme (TACE) and are therefore useful in the treatment of arthritis, tumor metastasis, tissue ulceration, abnormal wound healing, periodontal disease, graft rejection, insulin resistance, bone disease and HIV infection. In particular, the compounds of the invention provide enhanced levels of inhibition of the activity of TACE in vitro and in cellular assay and/or enhanced selectivity over MMP-1 and are thus particularly useful in the treatment of diseases mediated by TNF.

Also according to the present invention, there are provided processes for producing the compounds of the present invention, which processes comprise one of the following:

a) reacting a compound of formula




wherein  $n$ ,  $X$ ,  $Y$ ,  $A$ ,  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_8$ ,  $R_9$ ,  $R_{10}$ , and  $R_{11}$  are as defined above or a reactive derivative thereof, with a compound of formula

$R_{12}NHOH$

wherein  $R_{12}$  is as defined above, to give a compound of formula I;

or

b) deprotecting a compound of formula:




-18-

wherein  $n$ ,  $X$ ,  $Y$ ,  $A$ ,  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ , and  $R_{12}$ , are defined above, and  $R_{30}$  is a suitable protecting group such as *t*-butyl, benzyl, and trialkylsilyl, to give a corresponding compound of formula I

or

- 5 c) cleaving a resin supported hydroxamate derivative containing the group



wherein  $n$ ,  $X$ ,  $Y$ ,  $A$ ,  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_8$ ,  $R_9$ ,  $R_{10}$ , and  $R_{11}$ , are defined above to give a compound of formula I;

- 10 or

- d) resolving a mixture (e.g. racemate) of optically active isomers of a compound of formula I to isolate one enantiomer or diastereomer substantially free of the other enantiomer or diastereomers;

or

- 15 e) acidifying a basic compound of formula I with a pharmaceutically acceptable acid to give a pharmaceutically acceptable salt;

or

- f) converting a compound of formula I having a reactive substituent group or site to a compound of formula I having a different substituent group or site.

20

With regard to process a) the reaction can be carried out by processes known in the art, e.g. by reaction of an acid chloride or mixed anhydride reactive derivative with the compound of formula  $R_{12}NHOH$ .

Removal of protecting groups, as illustrated by process b) can be carried out by processes known in the art to provide the hydroxamic acid, -see for example Scheme 16 below.

- 5        Process c) may be carried using a strong acid such as TFA to cleave the hydroxamate from the resin.

10        With regard to process d) standard separation techniques may be used to isolate particular enantiomeric or diastereomeric forms. For example a racemic mixture may be converted to a mixture of optically active diastereoisomers by reaction with a single enantiomer of a 'resolving agent' (for example by diastereomeric salt formation or formation of a covalent bond). The resulting mixture of optically active diastereoisomers may be separated by standard techniques (e.g. crystallisation or chromatography) and individual optically active diastereoisomers  
15        then treated to remove the 'resolving agent' thereby releasing the single enantiomer of the compound of the invention. Chiral chromatography (using a chiral support, eluent or ion pairing agent) may also be used to separate enantiomeric mixtures directly.

- 20        The compounds of formula I may be isolated in the form of a salt of a pharmaceutically acceptable acid e.g. an organic or inorganic acid by treatment with an acid such as described above.

25        With regard to process e) compounds of formula I having a reactive substituent group such as hydroxy or amino or site such as -S- can be converted to other compounds of formula I in known manner, e.g. alcohol to ester or ether. Reactive sites such as a sulfur atom can be oxidized to SO or SO<sub>2</sub>. (e.g. as shown in Scheme 4 below). If necessary reactive substituent groups may be protected during the synthesis of compounds of formula I and removed as a last step -see Schemes 4  
30        and 17 below.

PROCESS OF THE INVENTION.


5 The compounds of the present invention may be conveniently prepared according to one of the general processes outlined below.

The compounds of the present invention, where  $n = 0$ ,  $X = O, S$  or  $NR^7$ , and  $R^8$  and  $R^9$  taken with the carbon atom to which they are attached, form a six membered heterocyclic ring containing  $N-R^7$ ,  $S$  or  $O$  and  $A = S, SO$  or  $SO_2$  may be prepared according to one of the general processes outlined below.

10 As outlined in scheme 1, the appropriately substituted mercaptan derivative was alkylated using  $\alpha$ -bromo acetic acid ester derivative in refluxing chloroform using  $N,N$ -diisopropylethylamine as base. The sulfide derivative thus obtained was reacted with appropriately substituted propargyl bromide derivative in refluxing acetone using  $K_2CO_3$  as base. In the case of  $X = -N-R^7$  the  $N$ -alkylation can be carried  
15 out in  $DMF/NaH$  at room temperature. The sulfide derivative thus obtained was oxidized using  $m$ -chloroperbenzoic acid in  $CH_2Cl_2$  or by using Oxone in methanol/water. The sulfone thus obtained can be converted to the corresponding piperidine derivative by reacting it with bis(2-chloroethyl) $N$ -substituted amine derivative.

-21-

## SCHEME 1


**SYNTHESIS:**

- a. Diisopropylethylamine/ CHCl<sub>3</sub>/ RT/ 3 Hr; b. K<sub>2</sub>CO<sub>3</sub>/ Acetone/ Propargyl bromide  
 5 derivative; c. Oxone/ MeOH:THF/THF/ RT; d. K<sub>2</sub>CO<sub>3</sub>/ 18-Crown-6/ (C<sub>4</sub>H<sub>9</sub>)<sub>4</sub>NBr/  
 Acetone/Bis-2-chloroethyl N-substituted amine derivative/ Reflux; e: NaOH/  
 THF:MeOH/RT and (COCl)<sub>2</sub>/ NH<sub>2</sub>OH.HCl/ Et<sub>3</sub>N/ THF/ DMF.

Bis-2-chloroethyl N-substituted amines can be prepared from the substituted  
 10 diethanolamine and thionyl chloride. (Scheme 2). The cyclic product obtained by the  
 above mentioned operation, can be hydrolyzed to carboxylic acid and subsequently  
 converted to the hydroxamic acid as outlined in scheme 1.

-22-

SCHEME 2




a: Diisopropylethylamine/ $R^7Br$ / $CHCl_3$ /Reflux; b:  $SOCl_2$ / $CH_2Cl_2$ /Reflux

- 5 The corresponding sulfides and sulfoxides can be prepared starting from the corresponding saturated heterocyclic carboxylic acid derivative. (Scheme-3). N-Boc protected isonipecotic acid can be lithiated using tert-butyllithium and the resulting anion was reacted with appropriately substituted disulfides. The sulfide derivative can be converted to hydroxamic acids by the procedure outlined above.

10

SCHEME:3




a: tert-Butyllithium/ $-78^\circ C$ /THF/Bis(4-but-2-ynoxyphenyl)disulfide; b:  $(COCl)_2$ / $NH_2OH \cdot HCl$ / $Et_3N$ /DMF/ $CH_2Cl_2$ ; c: 1.  $HCl$ /Dioxane; 2.  $R^7Br$ / $Et_3N$ ; d:  $MeOH$ /30%  $H_2O_2$

15



These sulfides subsequently can be converted to the sulfoxides using 30% hydrogen peroxide at room temperature. The required disulfides can be prepared from the appropriately substituted thiol and DMSO/HCl oxidation. This procedure can be applied to any saturated, fused or non-fused heterocyclic carboxylic acid derivative. (Scheme 4)

SCHEME:4




- a: *tert*-Butyllithium/ $-78^\circ\text{C}$ / THF/Bis(4-but-2-ynyloxyphenyl)disulfide; b:  $(\text{COCl})_2$ /  
 10  $\text{NH}_2\text{OH}\cdot\text{HCl}/\text{Et}_3\text{N}/\text{DMF}/\text{CH}_2\text{Cl}_2$ ; c:  $\text{CH}_2\text{Cl}_2$ / HCl/ MeOH /  $\text{R}^7\text{Br}$ /  $\text{Et}_3\text{N}$ ; d: MeOH/30%  
 $\text{H}_2\text{O}_2$ ; e: Oxone/MeOH/THF/Rt.

Alternatively, sulfone derivatives can also be lithiated and carbonylated using either dry ice or  $\text{CO}_2$  gas. (Scheme 5). The sulfone derivative can be a mono

-24-

heterocyclic, bicyclic, benzo fused or hetero aryl such as pyridyl, thienyl, furanyl, pyrazinyl, pyrimidyl, thiazolyl fused ring systems.


SCHEME:5



5 a: n-Butyllithium and quench with  $\text{CO}_2$ ; b:  $(\text{COCl})_2/\text{DMF}/\text{NH}_2\text{OH}\cdot\text{HCl}/\text{Et}_3\text{N}$

The oxygen analogue can be prepared (Scheme 6) from the appropriately substituted alkynyloxy-benzenesulfonyl acetic acid ethyl ester and 2-chloroethyl ether. The corresponding pyran derivative can be hydrolyzed to carboxylic acid, which can be converted to the hydroxamic acid derivative.


SCHEME 6



15 a: 2-Chloroethyl ether/  $\text{K}_2\text{CO}_3$ /18-Crown-6/ $n\text{-(C}_4\text{H}_9)_4\text{Br}$ /Acetone/ Reflux;  
 b: 10N. NaOH/THF/MeOH/RT;  
 c:  $(\text{COCl})_2/\text{DMF}/\text{NH}_2\text{OH}\cdot\text{HCl}/\text{Et}_3\text{N}$ .

The thiols used as intermediates for the synthesis of compounds of the invention can be made according to Scheme 7. Thus, sulfonic acid salts **1**, where  $\text{XR}_{\text{so}}$  is a hydroxy, thiol or substituted amino moiety may be alkylated with acetylenes **2**, where J is a suitable leaving group such as halogen mesylate, tosylate, or triflate to give **3**. Acetylenes **2** are commercially available or known compounds, or they may be synthesized by known methods by those skilled in the art. The sulfonic acid salts **3** may be converted into the corresponding sulfonyl chloride or other sulfonylating agent **4** by known methods, such as reaction with oxalyl chloride, phosphorus oxychloride or other reagent compatible with substituents  $\text{R}_1$ ,  $\text{R}_2$  and  $\text{R}_3$  and the acetylene. The sulfonyl chloride **4** can then be reduced to the corresponding thiol **5** using triphenylphosphine in a suitable solvent mixture such as dichloromethane/DMF at a temperature of between  $-20^\circ\text{C}$  and  $30^\circ\text{C}$ .

## SCHEME 7




- 5 Alternatively, disulfide 6 may be converted into di-acetylene 7 by reaction with compounds 2, followed by reduction of the disulfide bond to provide the desired thiols 5. Bisacetylenes 7 may also be converted into thiols 5 via sulfonyl chlorides 4.
- 10 Alkylation of the phenol, thiophenol, aniline or protected aniline 8 with 2 to give 9, followed by reaction with chlorosulfonic acid provides sulfonic acids 10 which are readily converted into 4 with oxalyl chloride or similar reagents and subsequently

-27-

reduced to thiols **5**. Thiophenols **11** are also precursors to **5** via protection of the thiol with a triphenylmethyl or other suitable protecting group, alkylation of XH, where X is O, N or S, and deprotection of the sulfur.

Scheme 8:



Compounds of the invention wherein X is N, O, S, SO or SO<sub>2</sub>, can be synthesized according to **Scheme 8** and **Scheme 9**. Alkylation of the para-disubstituted aryl **14**, or its protected equivalent, with acetylene **2** in the presence of a base such as potassium carbonate in a polar aprotic solvent such as acetone or DMF at a temperature of between 20°C and 120°C provides the mono-propargylic ether **15**. Those skilled in the art will recognize that protecting groups may be required to avoid undesirable side reactions and increase the yield of the reaction. The need and choice of protecting group for a particular reaction is known to those skilled in the art. Reaction of this compound with  $\alpha$ -propiolactone, or a substituted propiolactone derivative (wherein the substituents are defined as before), in the presence of a base such as potassium t-butoxide in a polar solvent, or solvent mixture, such as THF or DMF affords the carboxylic acid **16**. Conversion of carboxylic acid **16** into the corresponding hydroxamic acid, **17**, is accomplished via formation of an activated ester derivative such as an acid chloride or acid anhydride followed by reaction with hydroxylamine. It is understood by those skilled in the art that when A is sulfur, in **Scheme 8** and all relevant subsequent **Schemes**, the sulfur can be oxidized to the corresponding sulfoxide or sulfone at any stage after formation of the thioether, using a suitable oxidant such as oxone, air, m-chloroperbenzoic acid or hydrogen peroxide.


Compounds **18** are also accessible from the Michael addition of compound **15** to a cyclic acrylate ester, or substituted acrylate ester (substituents are defined as before), to provide **18**, in which R<sub>30</sub> is hydrogen or a suitable carboxylic acid protecting group. Deprotection of the ester moiety then provides carboxylic acid, which can be converted into the analogous hydroxamic acid. Similarly, Michael addition of mono-protected 1,4-disubstituted aryl **19**, where XR<sub>25</sub> is hydroxy or protected hydroxy, thiol or amine, gives compound **20**. Unmasking of the protecting group gives thiol, aniline or phenol **21** which can be alkylated with propargyl derivative **2** to provide **18**. Mono protected compound **19** can also be reacted with  $\beta$ -propiolactone to provide **22**. Alternatively, **22** can be deprotected followed by alkylation to give **16** and **17**.

-29-

Synthesis of compounds of the invention wherein X is N, O, S, SO or SO<sub>2</sub>, and the linker between the proximal heteroatom and the hydroxamic acid is a one or three carbon chain can be synthesized according to **Scheme 9**. Compound **19**, where XR<sub>25</sub> is hydroxy or protected hydroxy, thiol or amine, can react with ester **24** or lactone **24a**, in which R<sub>30</sub> is hydrogen or a suitable carboxylic acid protecting group, with an appropriately substituted leaving group such as halogen, tosylate, mesylate or triflate, to provide **25**. Unmasking of the heteroatom X of compound **25** then provides **26**, which may next be alkylated with propargylic derivative **2** to give acetylene-ester **27**. Ester **27** can be converted into the corresponding hydroxamic acid **28** through conversion of the ester into the carboxylic acid by acid or base hydrolysis, followed by conversion into the hydroxamic acid as described in **Scheme 1**. Alternatively, compound **15**, prepared as shown in **Scheme 8**, can be alkylated directly with ester **24** or lactone **24a** to give **27** and then **28**. Substituents on the carbon alpha to the hydroxamic are defined as before.

15

## SCHEME 9




Compounds of the invention wherein A is a methylene or substituted  
 5 methylene group, and X is oxygen, can be obtained according to **Scheme 10**. Esters  
 or carboxylic acids **29**, commercially available or known in the literature, can be  
 converted into the corresponding phenols, **30**. Alkylation of the phenol with acetylene  
 2 gives the propargylic ethers, **31**, which can be converted into the corresponding  
 carboxylic acids and thence the hydroxamic acids, **33**, as described in **Scheme 1**.  
 10 Substituents on the carbon alpha to the hydroxamic, are defined as before.




-31-

SCHEME 10



- 5 Compounds of the invention wherein A is -SO<sub>2</sub>-, and R<sub>8</sub> and R<sub>9</sub> are not hydrogen, are available starting from 4-fluorobenzenethiol **34** as shown in **Scheme 11**. Deprotonation of the thiol followed by reaction with  $\alpha$ -propiolactone, or an acrylate ester, or ester derivative **24**, and subsequent oxidation of the resulting thioether provides sulfone-acid **35**. Displacement of the 4-fluoro substituent of **35**, or
- 10 its corresponding ester, with propargyl derivative **36**, wherein X is N, O or S, then provides sulfone **16**. Compound **16** can be converted into the compounds of the invention according to **Scheme 1**. Fluoroaryl **35** can also react with a masked hydroxyl, thiol or amino group (HXR<sub>40</sub>, wherein R<sub>40</sub> is a suitable protecting group) in the presence of a base such as sodium hydride in a polar aprotic solvent such as DMF
- 15 to provide **36**. Deprotection of **36** followed by alkylation with acetylenic derivative **2** then gives **16**.


## SCHEME 11



5


Compounds of the invention wherein X is NH are also available starting from the appropriate commercially available nitro aryl compound **38** (Scheme 12). Thus, the anion of compound **38** can be used to alkylate  $\beta$ -propiolactone, or a substituted derivative, or a cyclic acrylate ester to provide **40** and **39** respectively. Reduction of the nitro group followed by alkylation of the resulting aniline then gives **16**. Compound **38** can also be alkylated with ester derivative **24** to afford nitro-ester **40**, followed by reduction to give the corresponding aniline, analogous to compound **26** of Scheme 9.

SCHEME 12



- 5 Compounds of the invention wherein R<sub>11</sub>, alpha to the hydroxamic acid, is a hydroxy group can be obtained via epoxides 41, as shown in **Scheme 13**. These epoxides are available through the oxidation of the corresponding acrylate esters or by the Darzens reaction of an alpha-halo ester with a ketone. Reaction of the epoxide with thiol, phenol or aniline 19 in the presence of base or Lewis acid catalyzed
- 10 epoxide ring opening, provides alpha-hydroxy ester 42. Deprotection of 42 followed by alkylation with propargyl derivative 2 gives 44. Conversion of the ester of 44 into the analogous hydroxamic acid as described in **Scheme 1** then provides 45. Compounds 45, wherein A is sulfur, may be converted into the analogous sulfoxides or sulfones through oxidation with hydrogen peroxide, air, Oxone or other suitable reagent at this point. Similarly, thiol, phenol or aniline 15 can be reacted with 41 to
- 15 give 44. The hydroxyl group of compound 43 can also be manipulated through its conversion into a suitable leaving group, such as halide or sulfonate ester, followed by displacement with various nucleophiles including amines to provide 44.


## SCHEME 13



5 Another route to  $\alpha$ -hydroxy hydroxamic acids of the invention is shown in **Scheme 14**. Compound **15** can be alkylated with alcohol **46** to give **47**. Oxidation of the alcohol, with or without concomitant oxidation of the thioether (for  $A = S$ ), gives the aldehyde **48**. Reaction of aldehyde **48** with trimethylsilyl cyanide or other suitable reagent then provides the cyanohydrin **49**. Hydrolysis of the nitrile **49** into the

-35-


corresponding carboxylic acid followed by conversion into the hydroxamic acid as described in **Scheme 1** gives **50**.

**SCHEME 14**

Compounds described in the present invention (from Example 30 to 63) were prepared as per the **Schemes 15** and **Scheme 16**. In scheme 15, the t-Boc-protected ethyl isonipecotate **51** was carefully alkylated using diiodomethane to yield the moniodo compound **52**. This was subsequently converted to different hydroxamic acid derivatives as depicted in **Scheme 15**. In scheme 16, the N-Boc group was selectively removed using TMSOTf/ 2,6-Lutidine. After the derivatisation of the nitrogen, the O-tBu was removed using TFA in methylene chloride.


-36-

Scheme 15



-37-


Scheme 16



- 5 Alternatively, compounds ( wherein A = SO<sub>2</sub> and n = 0) described in examples 64 to 74 and 80 were prepared as depicted in **Scheme 17**.

-38-

## Scheme 17





**Example 1****1-(4-Bromo-benzyl)-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperdine-4-carboxylic acid hydroxyamide**5    **Step 1:**

To a stirred solution of 4-mercapto phenol (12.6 g, 100 mmol) and N,N-diisopropylethylamine (13.0 g, 100 mmol) in chloroform (200 ml), ethyl bromoacetate (17.0 g, 100 mmol) in chloroform (30 ml) solution was added slowly at room temperature. After the addition was complete, the reaction mixture was  
10    refluxed for 1 hr and cooled to room temperature. The reaction mixture was washed well with water, dried over anhydrous  $\text{MgSO}_4$ ; filtered and concentrated. The oily product obtained was taken to next step without purification.

## Step 2:

15    A mixture of  $\text{K}_2\text{CO}_3$  (15 gm, excess), (4-hydroxy-phenylsulfanyl)-acetic acid ethyl ester (5 g, 23.6 mmol) and 1-bromo-2-butyne (9.34 g, 35.4 mmol) was refluxed with stirring for 8 hrs. The reaction mixture was then cooled to room temperature and filtered. The filtrate was concentrated and extracted with chloroform. The chloroform layer was washed with water, dried over anhydrous  $\text{MgSO}_4$ , filtered and  
20    concentrated. The product obtained was taken to next step with out purification. Yield 6.0 g (96%); yellow oil; MS: 264.0 EI ( $\text{M}^+\text{H}$ ).

## Step 3:

To a stirred solution of (4-but-2-ynyloxy-phenyl sulfanyl)-acetic acid ethyl  
25    ester (101 g, 380 mmol) in MeOH: THF (3:1) (1000 ml), Oxone (670.0 g, excess) in water (1000 ml) was added at room temperature. The reaction mixture was stirred at room temperature for 8 hrs. The reaction mixture was then diluted with chloroform (600 ml) and filtered. The organic layer was separated and washed once with a saturated solution of  $\text{NaHSO}_3$  (400 ml). The chloroform layer was washed well with  
30    water, dried and concentrated. The oily product was dissolved in MeOH (100 ml)

-40-

and hexane (600 ml) was added. The separated colorless solid was filtered and washed with hexane. Yield 108 g (96%); mp. 91 - 93°C; MS: 297 (M<sup>+</sup>H)<sup>+</sup>.

## Step 4:

5 A mixture of diethanolamine (22.5 g, 150 mmol), 4-bromobenzyl bromide (25 g, 100 mmol) and N,N-diisopropylethylamine (19.0 g, 150 mmol) was refluxed for 24 hrs in chloroform (500 ml) solution. The reaction mixture was then concentrated and the residue was extracted with chloroform. It was washed well with water, dried over anhydrous MgSO<sub>4</sub>, filtered and concentrated. The crude product obtained was taken  
10 to next step with out purification. Yield 33.6 g (99%); Yellow oil, MS: 273.8 (M+H)<sup>+</sup>.

## Step 5:

2-[(4-Bromobenzyl)-(2-hydroxy-ethyl)-amino]-ethanol (33.28 g, 122 mmol)  
15 was dissolved in methanolic hydrogen chloride (100 ml) at 0° C. Methanol was removed in vacuo and the hydrochloride salt was suspended in CH<sub>2</sub>Cl<sub>2</sub> (300 ml). To a stirred solution of the above mentioned suspension, thionyl chloride (30 g, excess) was added slowly at room temperature. The reaction mixture was brought to gentle reflux for 3 hrs. The reaction mixture was then concentrated and the (4-bromo-  
20 benzyl)-bis-(2-chloro-ethyl)-amine was used in the next step with out purification. Yield: 47 g (99%); brown solid; mp 125° C; MS: 309.8 (M+H)<sup>+</sup>.

## Step 6:

A stirred mixture of anhydrous K<sub>2</sub>CO<sub>3</sub> (10 g, excess), 18-crown-6 (1 g),  
25 tetrabutylammonium bromide (1.0 g), (4-but-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester (2.8 g, 9.46 mmol) and (4-bromo-benzyl)-bis-(2-chloro-ethyl)-amine (4.9 g, 14.2 mmol) in anhydrous acetone (200 ml) was refluxed for 24 hrs. The reaction mixture was then cooled and filtered and the filtrate was concentrated. The crude product was extracted with chloroform, washed well with water, dried and  
30 concentrated. The brown colored material was purified by column chromatography

-41-

on silica gel by eluting with 50% ethylacetate : hexane. Yield 1.36 g (27%); brown oil; MS: 534 (M+H)<sup>+</sup>

## Step 7:

5        1-(4-Bromo-benzyl)-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-(4-bromo-benzyl)-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (1.36 g, 2.54 mmol) dissolved in THF:methanol (100: 50 ml) and 10 N NaOH (15 ml). The reaction mixture was stirred at room temperature for 24 hrs. The reaction mixture was then  
10 concentrated and residue was cooled and neutralized with concentrated HCl. The separated solid was extracted with chloroform:methanol (3:1) (300 ml) and washed with water. The chloroform layer was dried and concentrated. The product was crystallized from methanol. Yield 800 mg (62%); off white solid; mp 197 °C; MS: 507.9 (M+H)<sup>+</sup>

15

## Step 8:

To a stirred solution of 1-(4-bromo-benzyl)-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid (750 mg, 1.5 mmol) and DMF (1 ml) in CH<sub>2</sub>Cl<sub>2</sub> (100 ml), oxalyl chloride (508 mg, 4.0 mmol) in methylene chloride (2 ml) was  
20 added dropwise at 0° C. After the addition, the reaction mixture was warmed to room temperature and stirred for 1 hr. The acid chloride thus formed was concentrated to remove excess oxalyl chloride and redissolved in CH<sub>2</sub>Cl<sub>2</sub> (30 ml). In a separate flask, hydroxylamine hydrochloride (690 mg, 10 mmol) was dissolved in DMF (10 ml) and triethylamine (10 g, 10 mmol) was added. The reaction mixture was further diluted  
25 with acetonitrile (25 ml) and stirred at 0° C. The acid chloride was slowly added into the hydroxylamine and after the addition was complete, the reaction mixture was brought to room temperature and stirred for 24 hrs. The reaction mixture was concentrated and the residue was extracted with chloroform, washed well with water and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The product was purified by silica gel column  
30 chromatography by eluting it with 10% methanol:ethyl acetate. 270 mg of 1-(4-bromo-benzyl)-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid

-42-

hydroxyamide was isolated as a hydrochloride salt, a white powder. Yield 52%; mp 153 °C; MS: 522.9 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 1.85 (t, J=2.04 Hz, 3H), 2.23 (m, 2H), 2.49 (m, 2H), 2.83 (m, 2H), 3.36 (m, 2H), 4.28 (s, 2H) 4.89 (d, J=2.2 Hz, 2H), 7.18 (d, J= 9 Hz, 2H), 7.47 (d, J=8.1 Hz, 2H), 7.68 (m, 4H), 9.37 (s, 1H), 10.25 (s, 1H)

### Example 2

#### 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid hydroxyamide

2-[(2-Hydroxy-ethyl)-(4-methoxy-benzyl)-amino]-ethanol was prepared according to the general method as outlined in Example 1 (Step 4). Starting from diethanolamine ( 10.5g, 100mmol). and 4-methoxy benzyl chloride ( 15.6g, 100 mmol). Yield 21g, ( 98%); yellow oil; MS: 226 (M+H)<sup>+</sup>

Bis-(2-chloro-ethyl)-(4-methoxy-benzyl)-amine was prepared according to the general method as outlined in Example 1 (Step 5). Starting from 2-[(2-hydroxy-ethyl)-(4-methoxy-benzyl)-amino]-ethanol (11.2 g, 50mmol). Yield 14g, ( 99%); dark brown low melting solid; MS: 263 (M+H)<sup>+</sup>

4-(4-but-2-ynyloxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in Example 1. Starting from (4-but-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester (2 g, 6.73 mmol) and bis-(2-chloro-ethyl)-(4-methoxy-benzyl)-amine (2.61 g, 8.75 mmol) and following the procedure as outlined in Example 1 (Step 6) 2.5 g of the product was isolated. Yield 2.5 g (77%); yellow oil; MS: 486 (M+H)<sup>+</sup>

4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-but-2-ynyloxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid ethyl ester (2.5 g, 5.15 mmol) dissolved in THF:methanol (3:1, 200 ml) and 10 N NaOH (15 ml). The resulting

-43-

reaction mixture was worked up as outlined in Example 1 (Step 7). Yield 1.26 g (54%); off white solid; mp 223 °C; MS: 458 (M+H)<sup>+</sup>

Starting from 4-(4-but-2-ynyloxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid (1 g, 2.19 mmol) and following the procedure as outlined in Example 1, (Step 8), 350 mg of 4-(4-but-2-ynyloxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a hydrochloride salt, an off white solid. Yield 31%; mp 162 °C; MS: 473 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 1.86 (t, J= 2.13 Hz, 3H), 2.23 (m, 2H), 2.49 (m, 2H), 2.73 (m, 2H), 3.39 (m, 2H), 3.77 (s, 3H), 4.21 (d, J= 4.26 Hz, 2H), 4.89 (d, J=2.28 Hz, 2H), 6.99 (d, J= 8.7 Hz, 2H), 7.17 (d, J=9 Hz, 2H), 7.43 (d, J=8.4 Hz, 2H), 7.68 (d, J= 9 Hz, 2H), 9.37 (s, 1H), 10.21 (s, 1H), 11.17 (s, 1H).

### Example 3

#### 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-chloro-benzyl)-piperidine-4-carboxylic acid hydroxyamide

2-[(4-chlorobenzyl)-(2-hydroxy-ethyl)-amino]-ethanol was prepared according to the general method as outlined in Example 1 (Step 4). Starting from diethanolamine (14.3 g, 95 mmol). and 4-chlorobenzyl chloride (10.2 g, 63 mmol). Yield 12.1 g, (84%); yellow oil; MS: 230 (M+H)<sup>+</sup>

(4-Chloro-benzyl)-bis-(2-chloro-ethyl)-amine was prepared according to the general method as outlined in Example 1 (Step 5). Starting from 2-[(4-chlorobenzyl)-(2-hydroxy-ethyl)-amino]-ethanol (12 g, 52.4 mmol). Yield 41.27 g, (90%); yellow powder; mp 115 °C; MS: 303 (M+H)<sup>+</sup>

4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-chloro-benzyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in Example 1 (Step 6). Starting from (4-but-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester (4 g, 13.5 mmol) and (4-chloro-benzyl)-bis-(2-chloro-ethyl)-amine (4.9 g, 16.2 mmol). Yield 3.5 g (53%); white crystals; MP 91.8°C; MS: 490 (M+H)<sup>+</sup>

4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-chloro-benzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-but-2-ynyloxy-benzenesulfonyl)-1-(4-chloro-benzyl)-piperidine-4-carboxylic acid ethyl ester (3.14 g, 6.42 mmol) dissolved in THF:methanol 3:1 (100 ml) and 10 N NaOH (10 ml). The resulting reaction mixture was worked up as outlined in Example 1 (Step 7). Yield 2.37 g (80%); white solid; mp 205 °C; MS: 461.9 (M+H)<sup>+</sup>

Starting from 4-(4-but-2-ynyloxy-benzenesulfonyl)-1-(4-chloro-benzyl)-piperidine-4-carboxylic acid (2.31 g, 5.01 mmol) and following the procedure as outlined in Example 1 (Step 8), 790 mg of 4-(4-but-2-ynyloxy-benzenesulfonyl)-1-(4-chloro-benzyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a hydrochloride salt, a yellow solid. Yield 31%; mp 130 °C; MS: 476.9 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 1.856 (s, 3H), 2.23 (m, 2H), 2.73-2.89 (m, 4H), 3.37 (d, 2H), 4.28 (m, 2H), 4.89 (d, 2H), 7.18 (d, J = 8.94 Hz, 2H), 7.54 (s, 4H), (d, J=8.88 Hz, 2H), 9.40 (s, 1H), 10.3 (s, 1H).

#### Example 4

##### 1-Benzyl-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

Bis-(2-Chloro-ethyl)-benzyl amine was prepared according to the general method as outlined in Example 1 (Step 5). Starting from N-benzyl-diethanolamine (164.6 g, 844 mmol). Yield 178.5 g (79%); brown solid; MS: 231.9 (M+H)<sup>+</sup>

1-Benzyl-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in Example 1 (Step 6). Starting from (4-but-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester (2 g, 6.73 mmol) and bis-(2-chloro-ethyl)-benzyl amine (2.3 g, 8.8 mmol). Yield 3.33 g (99%); yellow oil; MS: 455.9 (M+H)<sup>+</sup>

-45-

1-Benzyl-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-benzyl-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (3 g, 6.6 mmol) dissolved in THF:methanol (3:1 150 ml) and 10 N NaOH (15 ml). The resulting reaction mixture was worked up as outlined in Example 1 (Step 7). Yield 1.65 g (59%); off white powder; mp 191 °C; MS: 428 (M+H)<sup>+</sup>

Starting from 1-benzyl-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid (1.55 g, 3.63 mmol) and following the procedure as outlined in Example 1 (Step 8), 1.08 g of 1-benzyl-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a hydrochloride salt, an off white powder. Yield 62%; mp 175 °C; MS: 443 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 1.85 (t, J= 2.16 Hz, 3H), 2.25 (m, 2H), 2.49 (m, 4H), 2.77 (m, 2H), 4.28 (d, J=4.3 Hz, 2H), 4.89 (d, J= 2.28, 2H), 7.18 (m, 2H), 7.46 (m, 5 H), 7.73 (m, 2H), 9.36 (s, 1H), 10.27 (s, 1H), 11.08 (s, 1H).

### Example 5

#### 1-(4-Bromo-benzyl)-4-(4-pent-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

(4-Pent-2-ynyloxy-phenylsulfanyl)-acetic acid ethyl ester was prepared according to the general method as outlined in Example 1 (Step 2). Starting from (4-hydroxy-phenylsulfanyl)-acetic acid ethyl ester (5 g, 30 mmol) and 2-pentynyl chloride (3.7 g, 36.6 mmol) 7.15 g of the product isolated. Yield 7.15 g (86%); brown oil; MS: 278 EI (M+H)<sup>+</sup>

(4-Pent-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester was prepared according to the general method as outlined in Example 1 (Step 3). Starting from (4-pent-2-ynyloxy-phenylsulfanyl)-acetic acid ethyl ester (7.04 g, 25.3 mmol) and oxone (25 g) (4-Pent-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester was isolated. Yield 8 g (99%); yellow oil; MS: 310.9 (M+H)<sup>+</sup>

1-(4-Bromo-benzyl)-4-(4-pent-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in Example 1 (Step 6). Starting from (4-pent-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester (4 g, 12.9 mmol) and (4-bromo-benzyl)-bis-(2-chloro-ethyl)-amine (5.83 g, 16.8 mmol), 2.85 g of the product was isolated. Yield 2.85 g (31%); low melting white solid; MS: 549.9 (M+H)<sup>+</sup>

1-(4-Bromo-benzyl)-4-(4-pent-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-(4-bromo-benzyl)-4-(4-pent-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (2.64 g 4.8 mmol) dissolved in THF:methanol (100:50 ml) and 10 N NaOH (10 ml). The resulting reaction mixture was worked up as outlined in Example 1 (Step 7). Yield 1.6 g (65%); off white solid; mp 217 °C; MS: 521.9 (M+H)<sup>+</sup>

Starting from 1-(4-bromo-benzyl)-4-(4-pent-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid (1.55 g, 2.98 mmol) and following the procedure as outlined in Example 1 (Step 8), 200 mg of 1-(4-bromo-benzyl)-4-(4-pent-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, a yellow solid. Yield 12%; mp 62 °C; MS: 536.9 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 1.069 (t, J = 7.47 Hz, 3H), 2.26 (m, 2H), 2.49 (m, 2H), 2.73 (m, 2H), 2.89 (s, 2H), 3.40 (d, 2H), 4.26 (d, 2H), 4.9 (m, 2H) 7.18 (m, 2H), 7.48 (d, J = 8.4 Hz, 2H), 7.66 (m, 4H), 10.39 (s, 1H), 11.19 (s, 1H).

#### Example 6

##### 1-(4-Bromo-benzyl)-4-(4-oct-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

(4-Oct-2-ynyloxy-phenylsulfanyl)-acetic acid ethyl ester was prepared according to the general method as outlined in Example 1 (Step 2). Starting from (4-hydroxy-phenyl-sulfanyl)-acetic acid ethyl ester (5 g, 30 mmol) and 1-bromo-2-octyne (6.9 g, 36.6 mmol) 8.9 g of (4-oct-2-ynyloxy-phenylsulfanyl)-acetic acid ethyl ester was isolated. Yield 8.9 g (92%); yellow oil; MS: 320 EI (M+H)<sup>+</sup>



-47-

(4-Oct-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester was prepared according to the general method as outlined in Example 1 (Step 3). Starting from (4-oct-2-ynyloxy-phenylsulfanyl)-acetic acid ethyl ester (8.8 g, 27.5 mmol) 8.45 g of (4-oct-2-ynyloxy-phenylsulfanyl)-acetic acid ethyl ester was isolated. Yield 8.45 g (87%); yellow oil; MS: 352 EI (M+H)<sup>+</sup>

1-(4-Bromo-benzyl)-4-(4-oct-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in Example 1 (Step 6). Starting from (4-oct-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester (4 g, 11.4 mmol) and (4-bromo-benzyl)-bis-(2-chloro-ethyl)-amine (5.13 g, 14.8 mmol) 1.47 g of 1-(4-bromo-benzyl)-4-(4-oct-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was isolated. Yield 1.47 g (22%); yellow solid; MS: 591.9 (M+H)<sup>+</sup>

1-(4-Bromobenzyl)-4-(4-oct-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared starting from 1-(4-bromobenzyl)-4-(4-oct-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (1.36 g, 2.3 mmol) dissolved in THF:methanol (50:50 ml) and 10 N NaOH (10 ml). The resulting reaction mixture was worked up as outlined in Example 1 (Step 7). Yield 660 mg (51%); off white solid; mp 199 °C; MS: 562 (M+H)<sup>+</sup>

Starting from 1-(4-bromo-benzyl)-4-(4-oct-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid (570 mg, 1.01 mmol) and following the procedure as outlined in Example 1 (Step 8), 100 mg of 1-(4-bromo-benzyl)-4-(4-oct-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a hydrochloride salt, a white powder. Yield 17%; mp 140 °C; MS: 579 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 0.828 (t, J=7.14 Hz, 3H), 1.25 (m, 6H), 1.38 (m, 2H), 2.27 (m, 2H), 2.49 (m, 4H), 2.73 (m, 2H), 4.03 (m, 2H), 4.91 (s, 2H), 7.18 (d, J= 9 Hz, 2H), 7.47 (d, J= 8.1 Hz, 2H), 7.68 (m, 4H), 9.43 (s, 1H), 10.25 (s, 1H), 11.19 (s, 1H).

## Example 7

**4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-fluoro-benzyl)-piperidine-4-carboxylic acid hydroxyamide**

5        2-[(4-Fluoro-benzyl)-(2-hydroxy-ethyl)-amino]-ethanol was prepared according to the general method as outlined in Example 1 (Step 4). Starting from diethanolamine (15.7 g, 150 mmol), and 4-fluoro-benzyl chloride (14.4 g, 100 mmol) 20 g of the product was isolated. Yield 20 g, (93%); yellow oil; MS: 215 (M+H)<sup>+</sup>

10

(4-Fluoro-benzyl)-bis-(2-chloro-ethyl)-amine was prepared according to the general method as outlined in Example 1 (Step 5). Starting from 2-[(4-fluoro-benzyl)-(2-hydroxy-ethyl)-amino]-ethanol (23.6g, 110 mmol) 28 gms of the product was isolated. Yield 28 g, (96%); brown solid; mp 98-99 °C; MS: 251 (M+H)<sup>+</sup>

15

4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-fluoro-benzyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in Example 1 (Step 6). Starting from (4-but-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester (5 g, 16.9 mmol) and (4-fluoro-benzyl)-bis-(2-chloro-ethyl)-amine (5.8 g, 20.1 mmol) 5.3 g of the product was isolated. Yield 5.3 g (67%); Brown oil; MS: 474 (M+H)<sup>+</sup>

20

4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-fluoro-benzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-but-2-ynyloxy-benzenesulfonyl)-1-(4-fluoro-benzyl)-piperidine-4-carboxylic acid ethyl ester (9.5g, 20 mmol) dissolved in THF:methanol 3:1 (100 ml) and 10 N NaOH (20 ml). The resulting reaction mixture was worked up as outlined in Example 1 (Step 7). Yield 5.7 g (63%); white solid; mp 106-106 °C; MS: 447 (M+H)<sup>+</sup>

25

30        Starting from 4-(4-but-2-ynyloxy-benzenesulfonyl)-1-(4-fluoro-benzyl)-piperidine-4-carboxylic acid (5.7 g, 13 mmol) and following the procedure as

-49-

outlined in Example 1 (Step 8), 4.1 g of 4-(4-but-2-ynyloxy-benzenesulfonyl)-1-(4-fluoro-benzyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a HCl salt, a yellow solid. Yield: 64%; mp 162-4 °C; MS: 461 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 1.92 (s, 3H), 2.02-2.32 (m, 6H), 2.86 (m, 2H), 3.41 (d, 2H), 4.84 (d, 2H),  
5 7.01 (d, J = 8.94 Hz, 2H), 7.15 (d, J=8.88 Hz, 2H), 7.25 (d, J=9 Hz, 2H), 7.74 (d, J=9.0 Hz, 2H), 9.4-9.7 (bs, 1H).

### Example 8

#### 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-cyano-benzyl)-piperidine-4-carboxylic acid hydroxyamide

10 4-[[Bis-(2-hydroxyethyl)-amino]-methyl]benzonitrile was prepared according to the general method as outlined in Example 1 (Step 4) starting from diethanolamine (10.2 g, 97 mmol) and α-bromo-p-tolunitrile (15.8g, 81 mmol). Yield, (68%); white solid; mp 163 °C MS: 221.2 (M+H)<sup>+</sup>

15 4-[[Bis-(2-chloroethyl)-amino]-methyl]benzonitrile was prepared according to the general method as outlined in Example 1 (Step 5) starting from 4-[[bis-(2-hydroxyethyl)-amino]-methyl]benzonitrile (33.28 g, 122 mmol). Yield g, (%); brown solid; mp °C; MS: (M+H)<sup>+</sup>

20 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-cyano-benzyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in Example 1 (Step 6). Starting from 4-(4-but-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester (5.86g, 19.8 mmol) and 4-cyano-benzyl-bis-(2-chloro-ethyl)-amine  
25 (5.4g, 18 mmol) 4.7 g of the product was isolated. Yield (52%); amber oil; MS: 481.0 (M+H)<sup>+</sup>

30 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-cyano-benzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-cyano-benzyl)-piperidine-4-carboxylic acid ethyl ester(4 g, 8.3 mmol) dissolved in THF:Methanol (60: 30 ml) and 10 N NaOH (10 ml). The resulting reaction mixture

-50-

was worked up as outlined in Example 1 (Step 7). Yield 1.8g (48%); off white solid; MS: 441.9 (M+H)<sup>+</sup>

Starting from 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-cyano-benzyl)-  
5 piperidine-4-carboxylic acid (1.8g, 4 mmol) and following the procedure as outlined  
in Example 1 (Step 8), 0.20 g of 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-cyano-  
benzyl)-piperidine-4-carboxylic acid hydroxamide was isolated as a hydrochloride  
salt, white solid. Yield 20%; mp 109.6 °C; MS: 468.0 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz,  
DMSO-d<sub>6</sub>): δ 1.86 (m, 3H), 2.25 (m, 4H), 2.5 (m, 2H), 2.85 (d, 2H), 4.39 (s, 2H),  
10 4.88 (s, 2H), 7.15-7.19 (d, J=13.2, 2H), 7.67-7.70 (d, J=13.5, 2H), 7.78 (m, 2H),  
7.96-7.99 (d, J=9.6, 2H), 9.42 (s, 1H), 10.14 (s, 1H), 11.20 (s, 1H)

#### Example 9

##### 15 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-methyl-benzyl)-piperidine- 4-carboxylic acid hydroxamide

2-[(2-Hydroxy-ethyl)-(4-methyl-benzyl)-amino]-ethanol was prepared  
according to the general method as outlined in Example 1 (Step 4). Starting from  
diethanolamine (4.84g, 46 mmol) and 4-methylbenzyl bromide (8.5g, 46 mmol), 8.2  
20 g of the product was isolated. Yield, (85%); white solid; MS: 210.1 (M+H)<sup>+</sup>

4-Methyl-benzyl-bis-(2-chloro-ethyl)-amine was prepared according to the  
general method as outlined in example 1 (Step 5). Starting from 2-[(2-Hydroxy-  
ethyl)-(4-methyl-benzyl)-amino]-ethanol (6.0 g, 20 mmol) 5.2 g of the product was  
isolated. Yield: (84%); yellow solid; mp 145-147 °C; MS: 245.9 (M+H)<sup>+</sup>

25 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-methyl-benzyl)-piperidine-4-  
carboxylic acid ethyl ester was prepared according to the general method as outlined  
in Example 1 (Step 6). Starting from 4-(4-but -2-ynyloxy-benzenesulfonyl)-acetic  
30 acid ethyl ester (5.75g, 19.0 mmol) and 4-methyl-benzyl-bis-(2-chloro-ethyl)-amine  
(6.04, 208 mmol) 6.47 g of the product was isolate. Yield: (72%); amber oil; MS: 470  
(M+H)<sup>+</sup>

4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-methyl-benzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-methyl-benzyl)-piperidine-4-carboxylic acid ethyl ester (6.4 g, 13.6 mmol) dissolved in THF:Methanol (30: 20 ml) and 10 N NaOH (15 ml). The resulting reaction mixture was worked up as outlined in Example 1 (Step 7). Yield 2.3g (48%); off white solid; mp 213 °C MS: 441.9 (M+H)<sup>+</sup>

Starting from 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-methyl-benzyl)-piperidine-4-carboxylic acid (2.0g, 5.0 mmol) and following the procedure as outlined in Example 1 (Step 8), 3.6g of 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-methyl-benzyl)-piperidine-4-carboxylic acid hydroxamide was isolated as a HCl salt, off-white solid. Yield 1.2g (28%); mp 188 °C; MS: 457.0 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 1.86 (s, 3H), 2.27 (m, 2H), 2.50 (m, 4H), 2.64 (m, 2H), 4.23-4.24 (d, J=4.5, 2H), 4.89 (d, J=1.8, 2H), 7.16-7.19 (d, J=9 2H), 7.24-7.26 (d, J=7.5, 2H), 7.37-7.40 (d, J=8.1, 2H), 9.36 (s, 1H), 10.11 (s, 1H), 11.20 (s, 1H)

#### Example 10

##### 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(3,4-dichloro-benzyl)-piperidine-4-carboxylic acid hydroxamide

2-[(2-Hydroxy-ethyl)-(3,4-dichloro-benzyl)-amino]-ethanol was prepared according to the general method as outlined in Example 1 (Step 4). Starting from diethanolamine (4.84g, 46 mmol) and 3,4-dichlorobenzyl chloride (8.97g, 46 mmol), 9.4 g of the product was isolated. Yield, (78%); white solid; MS: 264.3 (M+H)<sup>+</sup>

3,4-Dichloro-benzyl-bis-(2-chloro-ethyl)-amine was prepared according to the general method as outlined in Example 1 (Step 5). Starting from 2-[(2-Hydroxy-ethyl)-(3,4-dichloro-benzyl)-amino]-ethanol (10.7 g, 41 mmol), 10.7 g of the product was isolated. Yield: (84%); yellow solid; mp 218-220 °C; MS: 301.8 (M+H)<sup>+</sup>

4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(3,4-dichloro-benzyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined

in Example 1 (Step 6). Starting from 4-(4-but -2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester (6.1g, 23 mmol) and 3,4-dichloro-benzyl-bis-(2-chloro-ethyl)-amine (8.6g, 25 mmol), 4.9 g of the product was isolated. Yield:(41%); amber oil; MS: 523.8 (M+H)<sup>+</sup>

5

4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(3,4-dichloro-benzyl)-piperidine-4-carboxylic acid was prepared starting from 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(3,4-dichloro-benzyl)-piperidine-4-carboxylic acid ethyl ester (8.6 g, 16.4 mmol) dissolved in THF:Methanol (40: 30 ml) and 10 N NaOH (15 ml). The resulting reaction mixture was worked up as outlined in Example 1 (Step 7). Yield 2.1g (38%); off white solid; mp 232 °C MS: 495.9 (M+H)<sup>+</sup>

10

Starting from 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(3,4-dichloro-benzyl)-piperidine-4-carboxylic acid (2.06g, 4.0 mmol) and following the procedure as outlined in Example 1 (Step 8), 1.2g of 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(3,4-dichloro-benzyl)-piperidine-4-carboxylic acid hydroxamide was isolated as a HCl salt, off-white solid. Yield 1.2g (56%); mp 213 °C; MS: 510.9 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 1.86 (s, 3H), 2.30 (m, 2H), 2.50 (m, 4H), 2.80 (m, 2H), 4.40 (s, 2H), 4.90 (s, 2H), 7.16-7.19 (d, J=9 2H), 7.51-7.54 (d, J=8.4, 2H), 7.66-7.69 (d, J=9.0, 2H), 7.75-7.86 (d, J=11.7, 2H), 7.88 (s, 1H), 9.38 (s, 1H), 10.44 (s, 1H), 11.19 (s, 1H).

15

20

### Example 11

#### 1-(4-Bromo-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide

25

Step 1:

(4-Prop-2-ynyloxy-phenylsulfanyl)-acetic acid ethyl ester was prepared according to the general method as outlined in Example 1 (Step 2). Starting from (4-hydroxy-phenylsulfanyl)-acetic acid ethyl ester (example 1, 1<sup>st</sup> paragraph) ( 2.12g, 10 mmol) and propargyl bromide ( 1.8g, 15mol) 2.4 g of the product was isolated. Yield: (96%); amber oil; MS: 251(M+H)<sup>+</sup>

30

## Step 2:

(4-Prop-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester was prepared according to the general method as outlined in Example 1 (Step 3). Starting from (4-prop-2-ynyloxy-phenyl sulfanyl)-acetic acid ethyl ester ( 2.5g, 10 mmol) 2.8 g of (4-  
5 Prop-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester was isolated. Yield (99%); brown oil ; MS: 283 (M+H)<sup>+</sup>

## Step 3:

1-(4-Bromo-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-  
10 carboxylic acid ethyl ester was prepared according to the general method as outlined in Example 1 (Step 6). Starting from (4-prop-2-ynyloxy-benzenesulfonyl)- acetic acid ethyl ester (21.62 g, 76.7 mmol) and (4-bromo-benzyl)-bis-(2-chloro-ethyl)-amine (31.9g, 92 mmol), 23 g of the ester derivative was isolated. Yiel: (58%); yellow oil; MS: 521.9 (M+H)<sup>+</sup>.

15

## Step 4:

1-(4-Bromo-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-  
carboxylic acid was prepared starting from 1-(4-bromo-benzyl)-4-(4-prop-2-ynyloxy-benzene-sulfonyl)-piperidine-4-carboxylic acid ethyl ester (5 g, 9.59 mmol) dissolved  
20 in THF:methanol (150:50 ml) and 10 N NaOH (15 ml). The resulting reaction mixture was worked up as outlined in Example 1( Step 7). Yield 3.4 g (72%); brown low melting solid; MS: 491.9 (M-H)<sup>-</sup>

## Step 5:

25 Starting from 1-(4-Bromo-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid (3 g, 6.1 mmol) and following the procedure as outlined in Example 1 (Step 8), 580 mg of 1-(4-bromo-benzyl)-4-(4-prop-2-ynyloxy-benzene-sulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as an HCl salt, off white powder. Yield 18%; mp 155 °C; MS: 508.8 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300  
30 MHz, DMSO-d<sub>6</sub>): δ 2.22 (m, 2H), 2.50 (m, 2H), 2.79 (m, 2H), 3.45 (m, 2H), 4.27 (m, 2H), 4.96 (d, J=2.3 Hz, 2H), 7.2 (d, J=9 Hz, 2H), 7.48 (m, 2H), 7.68 (m, 4H), 9.37 (s, 1H), 10.36 (s, 1H), 11.19 (s, 1H).

## Example 12

**1-(4-Bromo-benzyl)-4-[4-(4-piperidin-4-yl-but-2-ynyloxy)-benzenesulfonyl]-  
piperidine-4-carboxylic acid hydroxyamide**

5 To a stirred solution of piperidine (1.63 g, 19.2 mmol) diluted in dioxane (100 mL) acetic acid (5mL) was added. The reaction fumed and stirred for 5 minutes. 1-(4-bromo-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (5.0 g, 9.6 mmol), paraformaldehyde (0.29 g, 9.6 mmol) and the copper(I)chloride (0.35g) was added to the piperidine solution. The reaction turned  
10 green and was heated at reflux for 1 hour turned brown. It was then concentrated and diluted in ice water then brought to pH 8 with  $\text{NH}_4\text{OH}$  and extracted in  $\text{CHCl}_3$ . The organic layer was washed 4 times with water then dried over  $\text{Na}_2\text{SO}_4$  then concentrated. The product was purified by silica gel column chromatography by eluting it with 5% methanol: chloroform solution.

15 Yield 5.15 g (87%); brown oil; MS: 309.9 ( $\text{M}+2\text{H}$ )<sup>2+</sup>, 618.8 ( $\text{M}+\text{H}$ )<sup>+</sup>

1-(4-Bromo-benzyl)-4-[4-(4-piperidin-1-yl-but-2-ynyloxy)-benzenesulfonyl]-piperidine-4-carboxylic acid was prepared starting from 1-(4-bromo-benzyl)-4-[4-(4-piperidin-1-yl-but-2-ynyloxy)-benzenesulfonyl]-piperidine-4-carboxylic acid ethyl ester (4.64 g, 7.5 mmol) dissolved in THF:methanol (50:150 ml) and 10 N NaOH (20  
20 ml). The resulting reaction mixture was worked up as outlined in Example 1 (Step 7). Yield 3.35 g (76%); off white solid; mp 180 °C; MS: 295.9 ( $\text{M}+2\text{H}$ )<sup>2+</sup> 590.9 ( $\text{M}+\text{H}$ )<sup>+</sup>

Starting from 1-(4-bromo-benzyl)-4-[4-(4-piperidin-1-yl-but-2-ynyloxy)-benzene-sulfonyl]-piperidine-4-carboxylic acid (1.9 g, 3.2 mmol) and following the  
25 procedure as outlined in Example 1 (Step 8), 810 mg of 1-(4-bromo-benzyl)-4-[4-(4-piperidin-1-yl-but-2-ynyloxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide was isolated as a hydrochloride salt, a pale yellow solid. Yield 40%; mp 209 °C; MS: 303.4 ( $\text{M}+2\text{H}$ )<sup>2+</sup> 605.9 ( $\text{M}+\text{H}$ )<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ  
30 1.70 (m, 2H), 2.29 (m, 2H), 2.76 (m, 4H), 3.40 (m, 10H), 4.14 (s, 2H), 4.26 (2H),



-55-

7.24 (d, J= 9 Hz, 2H), 7.51 (d, J=8.4 Hz, 2H), 7.67 (m, 4H), 9.39 (s, 1H), 10.45 (s, 1H).

### Example 13

5     **1-(4-Bromo-benzyl)-4-[4-(4-morpholin-4-yl-but-2-ynyloxy)-benzenesulfonyl]-  
piperidine-4-carboxylic acid hydroxyamide**

To a stirred solution of morpholine (1.68 g, 19.2 mmol) diluted in dioxane (100 mL) acetic acid (5 mL) was added. The reaction fumed and stirred for 5 minutes. 1-(4-bromo-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-  
10 carboxylic acid ethyl ester (5.0 g, 9.6 mmol), paraformaldehyde (0.29 g, 9.6 mmol) and the copper(I)chloride (0.35g) was added to the piperidine solution. The reaction turned green and was heated at reflux for 1 hour turned brown. It was then concentrated and diluted in ice water then brought to pH 8 with  $\text{NH}_4\text{OH}$  and extracted in  $\text{CHCl}_3$ . The organic layer was washed 4 times with water then dried over  $\text{Na}_2\text{SO}_4$   
15 then concentrated. The product, 1-(4-Bromo-benzyl)-4-[4-(4-morpholin-4-yl-but-2-ynyloxy)-benzenesulfonyl]-piperidine-4-carboxylic acid ethyl ester was purified by silica gel column chromatography by eluting it with 5% methanol: chloroform solution. Yield 3.0 g (50%); colorless solid ; mp 110° C; MS: 311 ( $\text{M}+2\text{H}$ )<sup>2+</sup>, 621 ( $\text{M}+\text{H}$ )<sup>+</sup>

20

1-(4-Bromo-benzyl)-4-[4-(4-morpholin-4-yl-but-2-ynyloxy)-benzene-sulfonyl]-piperidine-4-carboxylic acid was prepared starting from 1-(4-bromo-benzyl)-4-[4-(4-morpholin-4-yl-but-2-ynyloxy)-benzenesulfonyl]-piperidine-4-  
carboxylic acid ethyl ester (2.87g, 4.6 mmol), dissolved in THF:methanol (3:1, 150  
25 ml) and 10 N NaOH (10 ml). The resulting reaction mixture was worked up as outlined in Example 1 (Step 7). Yield 2.26 g (83%); white powder; mp 198 °C; MS: 593.1 ( $\text{M}+\text{H}$ )<sup>+</sup>

Starting from 1-(4-bromo-benzyl)-4-[4-(4-morpholin-4-yl-but-2-ynyloxy)-  
30 benzenesulfonyl]-piperidine-4-carboxylic acid (2.1 g, 3.55 mmol) and following the procedure as outlined in example 1, 1.8 g of 1-(4-bromo- benzyl)-4-[4-(4-morpholin-

-56-

4-yl-but-2-ynyloxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide was isolated as a hydrochloride salt, a white solid. Yield 80%; mp 94 °C; MS: 304.4 (M+2H)<sup>2+</sup> 607.9 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 2.38 (m, 2H), 2.46 (m, 2H), 2.75 (m, 2H), 3.35 (m, 2H), 3.87, (m, 8H), 4.21 (s, 2H), 4.26 (s, 2H), 5.10 (s, 2H), 7.24 (d, J = 9 Hz, 2H), 7.51 (d, J = 8.4 Hz, 2H), 7.67 (m, 4H), 9.42 (s, 1H), 10.69 (s, 1H), 11.13 (s, 1H)

Examples of compound where A = S or S=O.

10

#### Example 14

##### 4-(4-But-2-ynyloxy-phenylsulfanyl)-4-hydroxycarbamoyl-piperidine-1-carboxylic acid tert-butyl ester

To a solution of triphenylphosphine (24.7g, 94.2 mmol) and dimethylformamide (0.6 mL) in dichloromethane (25 mL) was added a solution of 4-but-2-ynyloxy-phenylsulfonyl chloride (7.69g, 31.4 mmol) in dichloromethane dropwise over 30 min. After an additional 2h, 1N aqueous hydrochloric acid (20 mL) and water was added. The organic layer was separated and concentrated in vacuo. Aqueous sodium hydroxide (1N, 50 mL) was added and the solid removed by filtration. The aqueous phase was washed with diethyl ether (3x), treated with 1N aqueous hydrochloric acid (50 mL) and extracted with ether (3x). the combined organic extracts were dried over anhydrous magnesium sulfate and concentrated to give the thiol as an oil (3.77g). This material was dissolved in dimethylsulfoxide (40 mL) and concentrated hydrochloric acid was added (2 mL). After 18h, diethyl ether was added and the organic phased was washed with water (5x) and dried over anhydrous magnesium sulfate. Concentration in vacuo gave a yellow solid which was filtered through silica gel with hexane:ethyl acetate to give bis (4-but-2-ynyloxy phenyl) disulfide as a yellow solid (3.0g, 80%). <sup>1</sup>HNMR (CDCl<sub>3</sub>: 300MHz): 1.86 (s, -CH<sub>3</sub>, 3H), 4.63 (s, -CH<sub>2</sub>, 2H), 6.90 (d, ArH, 2H, J = 9 Hz), 7.40 (d, ArH, 2H, J = 9 Hz).

30

To a solution of N-BOC-isonipecotic acid (0.62g, 2.7 mmol) in tetrahydrofuran (20 mL) at -78 °C was added tert-butyllithium (3.4 mL, 1.7M in

-57-

hexane, 5.7 mmol). After 10 min at -78 °C the yellow solution was warmed to 0 °C in an ice bath. After 30 min the colorless solution was cooled to -78 °C whereupon bis (4-but-2-ynyloxy phenyl) disulfide (1.0 g, 2.8 mmol) was added as a solution in tetrahydrofuran (6 mL). The reaction mixture was allowed to warm to 25 °C. After 5 1.5h ethyl acetate was added followed by 6 mL of 1N aqueous hydrochloric acid in 20 mL of water. The organic layer was washed with water and brine, dried over anhydrous magnesium sulfate and concentrated in vacuo. Chromatography on silica gel (methanol/methylene chloride) gave the product (0.55g). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 1.38 (s, OtBu, 9H), 1.5 - 1.6 (m, CH<sub>2</sub>, 2H), 1.84 (s, CH<sub>3</sub>, 3H), 1.89 - 1.99 (m, CH<sub>2</sub>, 2H), 2.95 - 3.05 (m, CH<sub>2</sub>, 2H), 3.6 - 3.7 (m, CH<sub>2</sub>, 2H), 4.8 (s, CH<sub>2</sub>, 2H), 6.95 (d, 10 ArH, 2H, J = 9 Hz), 7.38 (d, ArH, 2H, J = 9 Hz).

Dimethylformamide (0.163 mL) was added to a solution of oxalyl chloride (1.06 mL of a 2.0M solution in dichloromethane) in dichloromethane (2 mL) at 0 °C. 15 After 15 min a solution of the acid in dimethylformamide (5 mL) was added and the reaction mixture was allowed to warm to room temperature. After 1h the reaction mixture was added to a mixture of hydroxylamine hydrochloride (0.737 g), triethylamine (2.22 mL), water (5.7 mL) and tetrahydrofuran (22.8 mL) that had been stirring at 0 °C for 15 min. The reaction was held at 0 °C for 18h then diluted with 20 dichloromethane and washed with saturated aqueous sodium bicarbonate (3X), then dried over potassium carbonate and concentrated in vacuo to give 480 mg of 4-(4-but-2-ynyloxy-phenylsulfanyl)-4-hydroxycarbonyl-piperidine-1-carboxylic acid tert-butyl ester. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 1.37 (s, OtBu, 9H), 1.5 - 1.6 (m, CH<sub>2</sub>, 2H), 1.84 (s, CH<sub>3</sub>, 3H), 1.9 - 2.0 (m, CH<sub>2</sub>, 2H), 3.05 - 3.15 (m, CH<sub>2</sub>, 2H), 3.5 - 3.6 (m, 25 CH<sub>2</sub>, 2H), 4.8 (s, CH<sub>2</sub>, 2H), 6.9 (d, ArH, 2H), 7.4 (d, ArH, 2H), 8.8 (s, NHOH, 1H), 10.7 (d, NHOH, 1H).

-58-

**Example 15****4-(4-But-2-ynyloxy-phenylsulfanyl)-piperidine-4-carboxylic acid hydroxyamide**

4-(4-But-2-ynyloxy-phenylsulfanyl)-4-hydroxycarbonyl-piperidine-1-carboxylic acid tert-butyl ester, prepared by the method outlined in Example 14 (Step 3) (0.175g, 0.4 mmol), was treated with 4N hydrochloric acid in dioxane (5 mL) at 25 °C for 1h 15 min. The reaction mixture was concentrated in vacuo, diethyl ether was added and the resulting precipitate isolated by filtration to give 4-(4-but-2-ynyloxy-phenylsulfanyl)-piperidine-4-carboxylic acid hydroxyamide as a white solid (0.12g).  
Electrospray Mass Spectroscopy:  $((M+H)^+ = 321)$

**Example 16****1-(4-Bromo-benzyl)-4-(4-but-2-ynyloxy-phenylsulfanyl)-piperidine-4-carboxylic acid hydroxyamide**

4-(4-But-2-ynyloxy-phenylsulfanyl)-piperidine-4-carboxylic acid hydroxyamide (prepared by the procedure outlined in example 15) (0.15g, 0.5 mmol) in methanol (5 mL) and dimethylformamide (2.5 mL) was treated with triethylamine (0.15 mL, 1.1 mmol) followed by 4-bromobenzylbromide (0.13g, 0.53 mmol). After 6h the solution was diluted with ethyl acetate, acidified to pH = 6 with 1N aqueous hydrochloric acid, washed sequentially with water, aqueous sodium bicarbonate and brine and dried over anhydrous sodium sulfate. Concentration in vacuo gave 1-(4-bromo-benzyl)-4-(4-but-2-ynyloxy-phenylsulfanyl)-piperidine-4-carboxylic acid hydroxyamide. <sup>1</sup>HNMR (DMSO-d<sub>6</sub>): 1.5 - 1.6 (m, CH<sub>2</sub>, 2H), 1.8 (s, CH<sub>3</sub>, 3H), 1.9 - 2.2 (m, CH<sub>2</sub>, 4H), 2.5 - 2.6 (m, CH<sub>2</sub>, 2H), 3.4 (s, CH<sub>2</sub>Ar, 2H), 4.75 (s, CH<sub>2</sub>, 2H), 6.9 (d, ArH, 2H), 7.2 (d, ArH, 2H), 7.3 (d, ArH, 2H), 7.5 (d, ArH, 2H), 8.8 (s, NHOH, 1H), 10.6 (d, NHOH, 1H). Electrospray Mass Spectroscopy:  $((M+H)^+ = 489/491)$

Examples of compounds, where n = 1 and A = S, S=O or SO<sub>2</sub>

**Example 17****4-(4-But-2-ynyloxy-phenylsulfanylmethyl)-tetrahydro-pyran-4-carboxylic acid hydroxyamide****4-But-2-ynyloxy-benzenesulfonic acid sodium salt**

5 To a solution of 52.35g (0.225 mol) of 4-hydroxybenzenesulfonate sodium salt in 1L of isopropanol and 225 mL of a 1.0N solution of sodium hydroxide was added 59.96g (0.45 mol) of 1-bromo-2-butyne. The resulting mixture was heated to 70° for 15h and then the isopropanol was removed by evaporation in vacuo. The resulting white precipitate was collected by filtration, washed with isopropanol and  
10 ether and dried in vacuo to give 56.0g (100%) of the butynyl ether as a white solid.

**4-But-2-ynyloxy-benzenesulfonyl chloride**

To a 0° solution of 43.8 mL (0.087 mol) of oxalyl chloride in 29 mL of dichloro-methane was dropwise added 6.77 mL (0.087 mol) of DMF followed by  
15 7.24g (0.029 mol) of 4-but-2-ynyloxy-benzenesulfonic acid sodium salt. The reaction mixture was stirred for 10 minutes at 0° then let warm to room temperature and stirred for 2 days. The reaction was then poured into ice and extracted with 150 mL of hexanes. The organics were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo to provide 6.23g (88%) of the sulfonyl chloride as  
20 a yellow solid; m.p. 63-65°C. EI Mass Spec: 243.9 (M<sup>+</sup>).

**But-2-ynyloxy-benzene**

To a solution of 6.14g (23.40 mmol) of triphenylphosphine dissolved in 100 mL of benzene and 50 mL of THF was added 1.75 mL (23.40 mmol) of 2-butyne-1-ol.  
25 After five minutes 2.00g (21.28 mmol) of the phenol, dissolved in 10 mL of THF, was added to the reaction followed by 3.69 mL (23.40 mmol) of diethyl azodicarboxylate. The resulting reaction mixture was stirred for 18h at room temperature and then concentrated in vacuo. The residue was chromatographed on silica gel eluting with ethyl acetate/hexanes (1:10) to provide 2.18g (70%) of the  
30 desired propargylic ether as a clear liquid. EI Mass Spec: 146.0 M<sup>+</sup>

-60-

## 4-But-2-ynyloxy-benzenesulfonyl chloride

To a solution of 0.146g (1.0 mmol) of the but-2-ynyloxy-benzene in 0.3 mL of dichloromethane in an acetone/ice bath under  $N_2$  was dropwise added a solution of 0.073 mL (1.1 mmol) of chlorosulfonic acid in 0.3 mL of dichloromethane. After the addition was complete, the ice bath was removed and the reaction was stirred at room temperature for 2h. To the reaction was then dropwise added 0.113 mL (1.3 mmol) of oxalyl chloride, followed by 0.015 mL DMF. The reaction was heated to reflux for 2h and then diluted with hexane and poured into ice water. The organic layer was washed with brine, dried over sodium sulfate, and concentrated in vacuo to provide 0.130g (53%) of the desired product as a light brown solid.

## 4-But-2-ynyloxy-benzenethiol

To a solution of 11.8g (.045 mol) of triphenylphosphine dissolved in 10 mL of dichloromethane and 0.3 mL of DMF was added 3.67g (.015 mol) of the 4-but-2-ynyloxy-benzenesulfonyl chloride, dissolved in 15 mL of dichloromethane and the resulting mixture was stirred for 2h at room temperature. After the addition of 5 mL of 1N HCl solution the reaction was stirred for 0.5h followed by the addition of 15 mL of brine. The organics were separated and concentrated in vacuo and the residue was diluted with ether and 2.5N sodium hydroxide solution. The resulting precipitate was filtered off and the aqueous layer was acidified to pH2 and extracted with ether. The combined organics were washed with brine, dried over  $Na_2SO_4$ , filtered through Magnesol® and concentrated in vacuo. The residue was chromatographed on silica gel eluting with hexanes/ether (4:1) to provide 1.13g (42%) of the thiol as a yellow oil. CI Mass Spec: 179 (M+H).

## 4-(4-But-2-ynyloxy-phenylsulfanylmethyl)-tetrahydro-pyran-4-carboxylic acid

To a solution of 0.112g (2.81 mmol) of 60% sodium hydride in 2 mL of THF, cooled to 0° C, was added a solution of 0.500g (2.81 mmol) of 4-but-2-ynyloxy-benzenethiol, dissolved in 3 mL of THF. The resulting mixture was stirred for 0.5h at room temperature, then cooled to 5° C, followed by the addition of 0.518g (3.65 mmol) of neat 2,7-dioxaspiro[3,5]nonane-1-one while keeping the reaction

-61-

temperature below 10° C. The reaction was allowed to warm to room temperature and stirred for an additional 0.5h and then quenched with 3 mL of 3N HCl solution and 3 mL of water. The resulting mixture was extracted with dichloromethane and the combined organics were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered through a plug of silica gel and concentrated in vacuo. The residue was triturated with hexanes and acetonitrile and filtered to give 0.72g of the carboxylic acid as a semi-solid. Electrospray Mass Spec: 319 (M-H)<sup>-</sup>

4-(4-But-2-ynyloxy-phenylsulfanylmethyl)-tetrahydro-pyran-4-carboxylic acid hydroxyamide

To a 0° C solution of 0.74g (2.31 mmol) of the product of 4-(4-but-2-ynyloxy-phenylsulfanylmethyl)-tetrahydro-pyran-4-carboxylic acid, dissolved in 7 mL of dichloromethane and 0.175 mL of DMF was added 1.27 mL (2.54 mmol) of a 2M solution of oxalyl chloride. The reaction was warmed to room temperature and stirred for 2h and then recooled to 0° C. A mixture of 0.875 mL (14.2 mmol) of a 50% hydroxylamine solution, 5.0 mL of THF and 2.0 mL of t-butanol were then added to the reaction. The reaction was stirred at room temperature for 1h and then concentrated in vacuo. The residue was extracted with dichloromethane and the combined organics were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The residue was chromatographed on silica gel eluting with dichloromethane/methanol (92:8) to provide 0.212g of the sulfide-hydroxamic acid as a white solid; m.p.135-137°C. Electrospray Mass Spec: 336 (M+H)<sup>+</sup>

Example 18

4-(4-But-2-ynyloxy-benzenesulfonylmethyl)-tetrahydro-pyran-4-carboxylic acid hydroxyamide

To a 0° C solution of 0.186g (0.56 mmol) of the product of 4-(4-but-2-ynyloxy-phenylsulfanylmethyl)-tetrahydro-pyran-4-carboxylic acid hydroxyamide, dissolved in 1.2 mL of THF and 4.8 mL of methanol was dropwise added a solution of 0.619g (1.008 mmol) of Oxone® in 3 mL of water, while keeping the temperature below 20° C. After the addition was complete the reaction was stirred at room

-62-

temperature for 3h. The reaction mixture was then poured into a cooled solution of 2.5 mL of toluene and 5 mL of ethyl acetate and the precipitate was filtered off. The filtrate was extracted with ethyl acetate/toluene and the combined organic layers were washed with water, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated in vacuo. The residue was  
5 triturated with ethyl acetate/toluene (5:2), filtered and dried in vacuo to provide 0.12g (55%) of the sulfone-hydroxamic acid as a white solid; m.p. 184-185° C. Electrospray Mass Spec: 368 (M+H)<sup>+</sup>

#### Example 19

##### 10 4-(4-But-2-ynyloxy-benzenesulfinylmethyl)-tetrahydro-pyran-4-carboxylic acid hydroxyamide

To a 0°C solution of 0.288g (0.80 mmol) of the product of 4-(4-but-2-ynyloxy-benzenesulfanylmethyl)-tetrahydro-pyran-4-carboxylic acid hydroxyamide dissolved in 20 mL of methanol was added 7.0 mL of 30% hydrogen peroxide  
15 solution. The reaction was allowed to warm to room temperature and stirred for 24h. The reaction mixture was then recooled to 0°C, quenched with saturated Na<sub>2</sub>SO<sub>3</sub> and concentrated in vacuo. The residue was diluted with water and dichloromethane. The organics were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The residue was chromatographed on silica gel eluting with  
20 dichloromethane/methanol (95:5) to provide 0.050g of the sulfoxide as a white solid. Electrospray Mass Spec: 351.9 (M+H)<sup>+</sup>

#### Example 20

##### 25 4-{{[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxytetrahydro-2H-pyran-4-carboxamide}

Step 1:

Ethyl 4-{{[4-(2-butynyloxy)phenyl]sulfonyl} tetrahydro-2H-pyran-4-carboxylate

(4-but-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester (10 g, 33.8 mmol) was added to a stirring solution of potassium carbonate (12 g), 18-crown-6 (0.5 g), 2-chloroethyl ether (4.75 ml, 40.5 mmol), and tetrabutyl ammonium bromide (0.5 g) in  
30 methyl ethyl ketone (200 ml). The mixture was heated at reflux overnight before the



-63-

salts were filtered off and the filtrate was concentrated. The residue was dissolved in chloroform and washed with water. The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated. The compound was isolated using silica-gel column chromatography by eluting it with 20% ethyl acetate: hexane solution. Ethyl 4-{{4-(2-butynyloxy)phenyl}sulfonyl} tetrahydro-2H-pyran-4-carboxylate was isolated as a yellow oil (10.06 g). Yield 80%; MS: 367.2 (M+H)<sup>+</sup>

4-{{4-(2-butynyloxy)phenyl}sulfonyl} tetrahydro-2H-pyran-4-carboxylic acid was prepared according to the general method as outlined in example 1 (step 7), starting from ethyl 4-{{4-(2-butynyloxy)phenyl}sulfonyl} tetrahydro-2H-pyran-4-carboxylate (10 g, 27.3 mmol); 2.7 g white solid. mp: 197 °C; Yield 30%; MS: 337.2 (M-H)<sup>-</sup>

Starting from a crude mixture of 4-{{4-(2-butynyloxy) phenyl} sulfonyl} tetrahydro-2H-pyran-4-carboxylic acid (2.59 g, 7.66 mmol), and following the procedure as outlined in Example 1 (step 8), 1.51 g of 4-{{4-(2-butynyloxy) phenyl} sulfonyl}-N-hydroxytetrahydro-2H-pyran-4-carboxamide was isolated as off white crystals. Mp: 210 °C; Yield: 58%; MS: 354.2 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  1.85 (t, J=2.28 Hz, 3H), 1.92 (m, 2H), 2.20 (d, J=13.1 Hz, 2H), 3.15 (t, J=11.52, 2H), 3.86 (d of d, 2H), 4.88 (d, J=2.34 Hz, 2H), 7.16 (d, J=8.7 Hz, 2H), 7.66 (d, J=8.91 Hz, 2H), 9.16 (s, 1H), 11 (s, 1H).

### Example 21

#### 1-benzyl-4-{{3-(2-butynyloxy)phenyl}sulfonyl}-N-hydroxy-4-piperidine carboxamide

Ethyl [(3-hydroxyphenyl) sulfanyl] acetate was prepared according to the general method as outlined in example 1 (step 1), starting from ethyl bromoacetate (7.95 g, 47.6 mmol) and 3-hydroxythiophenol (7.95 g, 47.6 mmol); 4.21 g yellow oil. Yield 41%; MS: 211.2 (M-H)<sup>-</sup>

Ethyl{[3-(2-butyloxy)phenyl]sulfanyl}acetate was prepared according to the general method as outlined in example 1 (step 2), starting from ethyl [(3-hydroxyphenyl)sulfanyl]acetate (3.87g, 18.3 mmol) and 4-bromo-2-butyne (2.66 g, 20 mmol); 5.16 g yellow oil. Yield 100%; MS(EI): 264.1 (M+H)<sup>+</sup>

5

Ethyl{[3-(2-butyloxy)phenyl]sulfonyl}acetate was prepared according to the general method as outlined in example 1 (step 3), starting from ethyl{[3-(2-butyloxy)phenyl]sulfanyl}acetate (5g, 18.9 mmol) and oxone (23.3 g, 37.9 mmol); 6.19 g yellow oil. Yield 100%; MS(EI): 296.1 (M+H)<sup>+</sup>

10

Ethyl 1-benzyl-4-{[3-(2-butyloxy)phenyl]sulfonyl}-4-piperidine carboxylate was prepared according to the general method as outlined in example 1 (step 6), starting from ethyl{[3-(2-butyloxy)phenyl]sulfonyl}acetate (3 g, 10.1 mmol) and Benzyl-bis- (2-chloro-ethyl) amine hydrochloride (2.88 g, 10.7 mmol); 2.91 g yellow oil. Yield 63%; MS: 456.3 (M+H)<sup>+</sup>

15

1-benzyl-4-{[3-(2-butyloxy)phenyl]sulfonyl}-4-piperidine carboxylic acid was prepared according to the general method as outlined in example 1 (step 7), starting from ethyl 1-benzyl-4-{[3-(2-butyloxy)phenyl]sulfonyl}-4-piperidine carboxylate (2.9 g, 6.37 mmol); 1.10 g off white powder. mp: 171 °C; Yield 40%; MS: 428.4 (M+H)<sup>+</sup>

20

Starting from 1-benzyl-4-{[3-(2-butyloxy)phenyl]sulfonyl}-4-piperidine carboxylic acid (1 g, 2.34 mmol), and following the procedure as outlined in Example 1 (step 8), 460 mg of 1-benzyl-4-{[3-(2-butyloxy)phenyl]sulfonyl}-N-hydroxy-4-piperidine carboxamide was isolated as an off white solid. mp: 91.4°C; Yield: 41%; MS: 443.4 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 1.83 (t, 3H), 2.23-2.27 (m, 2H), 2.73-2.89 (m, 2H), 3.29 (m, 2H), 3.68 (q, 2H), 4.31 (m, 1H), 4.39 (d, J=5 Hz, 1H), 4.85 (d, J=2.25, 2H), 7.25-7.61 (m, 9H), 9.1 (s, 1H), 11.2 (s, 1H).

25

-65-

**Example 22****4-{{4-(2-butynyloxy)phenyl}sulfonyl}-N-hydroxy-1-isopropyl-4-piperidine  
carboxamide**

- Ethyl 4-{{4-(2-butynyloxy)phenyl}sulfonyl}-1-isopropyl-4-piperidine  
5 carboxylate was prepared according to the general method as outlined in example 1  
(step 6), starting from (4-but-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester (6g,  
20.3 mmol) and isopropyl [bis(2-chloroethyl)] amine hydrochloride (4.88 g, 22.3  
mmol); 5.28 g brown oil. Yield 64%; MS: 408.2 (M+H)<sup>+</sup>
- 10 4-{{4-(2-butynyloxy)phenyl}sulfonyl}-1-isopropyl-4-piperidine carboxylic acid  
was prepared according to the general method as outlined in example 1 (step7),  
starting from ethyl 4-{{4-(2-butynyloxy)phenyl}sulfonyl}-1-isopropyl-4-piperidine  
carboxylate (5.25 g, 13 mmol); 2.06 g yellow solid. mp: 233 °C; Yield 42%; MS:  
380.1 (M+H)<sup>+</sup>
- 15 Starting from 4-{{4-(2-butynyloxy)phenyl}sulfonyl}-1-isopropyl-4-piperidine  
carboxylic acid (1.9 g, 5 mmol), and following the procedure as outlined in Example 1  
(step 8), 107 mg of 4-{{4-(2-butynyloxy)phenyl}sulfonyl}-N-hydroxy-1-isopropyl-4-  
piperidine carboxamide was isolated as a brown solid. mp: 105°C; Yield: 5%; MS:  
395.2 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 1.2 (m, 6H), 1.85 (t, 3H), 2.27 (m,  
20 2H), 2.73 (m, 2H), 3.06 (m, 2H), 3.52 (m, 2H), 3.57 (m, 1H), 4.89 (m, 2H), 7.19 (m,  
2H), 7.71 (m, 2H), 9.3 (s, 1H), 11.4 (s, 1H).

**Example 23****4-{{4-(2-butynyloxy)phenyl}sulfonyl}-N-hydroxy-1-(3-pyridinylmethyl)-4-  
25 piperidine carboxamide**

- Ethyl 4-{{4-(2-butynyloxy)phenyl}sulfonyl}-1-(3-pyridinylmethyl)-4-piperidine  
carboxylate was prepared according to the general method as outlined in example 1  
(step 6), starting from (4-but-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester (4g,  
16.9 mmol) and 3-pyridyl methyl [bis(2-chloroethyl)] amine hydrochloride (4.18 g,  
30 18.6 mmol); 370 mg brown oil. Yield 5%; MS: 457.4 (M+H)<sup>+</sup>

4-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-(3-pyridinylmethyl)-4-piperidine  
carboxylic acid was prepared according to the general method as outlined in example 1  
(step 6), starting from ethyl 4-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-(3-pyridinyl-  
methyl)-4-piperidine carboxylate (320 mg, 0.7 mmol); 150 mg yellow solid. Yield  
5 50%; MS: 429.2 (M+H)<sup>+</sup>

Starting from 4-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-(3-pyridinyl methyl)-4-  
piperidine carboxylic acid (860 mg, 2 mmol), and following the procedure as outlined  
in Example 1 (step 8), 800 mg of 4-{{[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxy-1-  
10 (3-pyridinylmethyl)-4-piperidine carboxamide was isolated as a white solid. mp:  
115°C; Yield: 84%; MS: 444.1 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 1.86 (t,  
J=1.98 Hz, 3H), 2.32 (m, 2H), 2.46 (s, 2H), 2.84 (m, 2H), 3.46 (d, J=12 Hz, 2H),  
4.45 (s, 2H), 4.89 (d, 2.1 Hz, 2H), 7.17 (d, J=8.9 Hz, 2H), 7.68 (d, J=8.85 Hz, 2H),  
7.9 (t, J=5.6 Hz, 1H), 8.0 (s, 1H), 8.51 (d, J=7.9 Hz, 1H), 8.87 (d, J= 4.6 Hz, 1H),  
15 8.99 (s, 1H), 11.4 (s, 1H).

#### Example 24

##### 3-{{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-ethyl-N-hydroxy-3- piperidinecarboxamide

20 Step 1: Piperidine-1,3-dicarboxylic acid 1-tert-butyl 3-ethyl ester

To a stirred solution of ethyl nipecotate (5.1g, 33 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (75 ml) and  
triethylamine (3.7g, 36 mmol) was added portionwise di-t-butylidicarbonate (7.1g, 33  
mmol). The reaction mixture was stirred at room temperature for 18 h, quenched with  
ice water and extracted with chloroform. The organic layer was dried over sodium  
25 sulfate, filtered, concentrated and chromatographed on a silica-gel column with 20:80  
ethyl acetate:hexane. Piperidine 1,3dicarboxylic acid 1-tert-butyl ester-3-ethyl ester  
was isolated as a waxy solid. Yield 6.86 g ( 82%). MS (ES): m/z 258.2 (M+H)<sup>+</sup>.

-67-

Step 2: 1-(tert-Butyl) 3-ethyl 3-{{[4-2-butynyloxy]phenyl}sulfonyl}-1,3-piperidine dicarboxylate

To a stirred solution of diisopropylamine (7.2g, 28 mmol) in THF (25 ml) at -78° C was added n-butyllithium (1.6m solution in hexanes, 19.0 ml, 30.8 mmol). The mixture was stirred for 30 min at 0°C. The mixture was then cooled to -78° C and piperidine -1,3-dicarboxylic acid 1-tert-butyl ester 3-ethyl ester (5.3g, 28 mmol) in THF (20 ml) was added slowly. The reaction mixture was stirred for 30 min then 4-but-2-ynyloxy-benzenesulfonyl fluoride (6.4g, 28 mmol) in THF (15 ml) was added slowly. The reaction was warmed to room temperature and after 4 hrs quenched with ice water and extracted with chloroform. The organic layer was dried over sodium sulfate, filtered, concentrated and chromatographed on a silica-gel column with 20% ethyl acetate:hexane to afford 1-(tert-Butyl) 3-ethyl 3-{{[4-2-butynyloxy]phenyl}sulfonyl}-1,3-piperidine dicarboxylate as a white solid. Yield 9.8 g (76%); mp 103.4°C; MS (ES): m/z 466.4 (M+H)+. <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 1.07 (t, 3H), 1.34 (s, 9H), 3.31 (s, 3H), 3.84 (m, 2H), 4.00 (m, 4H), 4.53 (d, 2H), 4.91 (m, 4H), 7.22 (d, 2H), 7.71 (d, 2H).

Step 3: To a stirred solution of 1-(tert-Butyl) 3-ethyl 3-{{[4-2-butynyloxy]phenyl}sulfonyl}-1,3-piperidine dicarboxylate (5.45g, 11.7 mmol) in methylene chloride (25 ml) at 0° C was added a saturated solution of hydrogen chloride in methylene chloride (25 ml). After 5 hours the solution was concentrated to afford ethyl 3-{{[4-(2-butynyloxy)phenyl}sulfonyl}-3-piperidinecarboxylate hydrogen chloride and is stored under nitrogen. White hygroscopic solid; Yield 3.47g (74%); MS (ES): m/z 366.2 (M+H)<sup>+</sup>

Step 4: (Ethyl 3-{{[4-(2-butynyloxy)phenyl}sulfonyl}-1-ethyl-3-piperidine-carboxylate)

3-{{[4-(2-Butynyloxy)phenyl}sulfonyl}-3-piperidinecarboxylate hydrogen chloride (2.97g, 8.0 mmol), ethyl iodide (1.28g, 8 mmol) and dry powdered potassium carbonate (3.8g) in dry acetone (60 ml) was heated to reflux for 18 hours. The mixture was allowed to cool and the potassium salts were filtered and concentrated. The

-68-

residue was extracted with chloroform and washed with H<sub>2</sub>O, dried over sodium sulfate and concentrated to afford ethyl 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-ethyl-3-piperidinecarboxylate. This product was used without further purification. Amber gum, yield 3.47 g (99%); MS (ES): m/z 394 (M+H)<sup>+</sup>.

5

Step 5 : 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-ethyl-3-piperidinecarboxylic acid  
3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-ethyl-3-piperidinecarboxylic acid was prepared starting from ethyl 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-ethyl-3-piperidinecarboxylate (3.2g, 8.0 mmol) dissolved in THF:Methanol (15:25 ml) and NaOH (15 ml). The resulting reaction mixture was worked up as outlined in example 1 (step 7). Yield 2.11g (71%), white solid; mp 159.2° C; MS (ES): m/z 366.3 (M+H) +

10

Step 6: 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-ethyl-N-hydroxy-3-piperidinecarboxamide

15 Starting from 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-ethyl-3-piperidinecarboxylic acid (2.0g, 5.5 mmol) and following the procedure as outlined in example 1 (step 8), 0.193g of 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-ethyl-N-hydroxy-3-piperidinecarboxamide hydrogen chloride was isolated as a white solid. Yield 10%; mp 190.3° C; MS (ES): m/z: 405.3 (M+H)<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 1.18 (m, 3H), 1.97 (m, 2H), 2.55 (m, 2H), 3.21 (m, 5H), 3.52 (s, 3H), 3.82 (d, 1H), 4.91 (m, 2H), 7.19 (d, 2H), 7.51 (s, 5H), 8.67 (s, 1H), 9.48 (s, 1H).

20

### Example 25

3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-(4-chlorobenzyl)-N-hydroxy-3-piperidinecarboxamide

25

Step 1: Ethyl 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-(4-chlorobenzyl)-3-piperidinecarboxylate

Starting from ethyl 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-3-piperidinecarboxylate hydrogen chloride (1.1g, 2.7 mmol) and 4-chlorobenzyl chloride (0.485, 3.0 mmol) in dry acetone (50 ml) and following the procedure outlined in example 24, (step 4), Ethyl 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-(4-chlorobenzyl)-3-piperidine-

30

-69-

carboxylate was isolated as a brown oil. This product was taken to the next step without further purification. Yield 1.66g (99%); MS (ES): m/z: 491.3 (M+H)<sup>+</sup>

Step 2: 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-(4-chlorobenzyl)-3-piperidine-  
 5 carboxylic acid  
 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-(4-chlorobenzyl)-1-3-piperidinecarboxylic acid  
 was prepared starting from ethyl 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-(4-chloro-  
 benzyl)-3-piperidinecarboxylate (1.64g, 3.3 mmol) dissolved in THF:Methanol (15:50  
 ml) and NaOH (15 ml). The resulting reaction mixture was worked up as outlined in  
 10 example 1 (step 7); Yield 1.11g (75%), white solid; mp 115.2° C; MS (ES): m/z 462.1  
 (M+H)<sup>+</sup>

Step 3: 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-(4-chlorobenzyl)-N-hydroxy-  
 3-piperidinecarboxamide  
 15 Starting from 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-(4-chlorobenzyl)-3-  
 piperidinecarboxylic acid (1.1g, 2.4 mmol) and following the procedure as  
 outlined in example 1, (step 8), 0.48g of 3-{{[4-(2-butynyloxy)phenyl]sulfonyl}-1-(4-  
 chlorobenzyl)-3-N-hydroxy-3-piperidinecarboxamide hydrogen chloride was isolated  
 as a white solid. Yield 43%; mp 124.4° C; MS (ES): m/z: 477.1 (M+H)<sup>+</sup>; <sup>1</sup>H NMR  
 20 (300 MHz, DMSO-d<sub>6</sub>): δ 2.0 (m, 2H), 3.39 (m, 5H), 4.27 (d, 2H), 4.89 (m, 2H), 7.14  
 (d, 2H), 7.15 (m, 4H), 7.61 (d, 2H), 8.95 (s, 1H), 9.46 (s, 1H).

### Example 26

4-{{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-  
 25 piperidine-4-carboxylic acid hydroxyamide.

A mixture of diethanolamine (2.1 g, 20 mmol), 4-(2-piperidin-1-yl-ethoxy)-benzyl  
 chloride (5.9 g, 20 mmol) and K<sub>2</sub>CO<sub>3</sub> (10 g, excess) was refluxed in acetone (100 ml)  
 for 24 hrs. At the end, reaction mixture was cooled to room temperature and filtered.  
 It was concentrated to dryness and redissolved in toluene (200 ml) and thionyl  
 30 chloride (6.75 g, 50 mmol). It was heated to 80° C for 1 hr and the separated brown  
 solid, bis-(2-chloro-ethyl)-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-amine was filtered

and dried. The crude product was taken to next step with out purification. Yield: 7.0 g, (89%).

4-{{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid ethylester was prepared according to the general method as outlined in example 1 (step 6), starting from ethyl{{[4-(2-butynyloxy)-phenyl]sulfonyl}acetate (2.9 g, 10.0 mmol) and bis-(2-chloro-ethyl)-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-amine dihydrochloride (4.3 g, 10 mmol), 2.8 g of product (brown oil) was isolated.. Yield 48%; MS: 583 (M+H)<sup>+</sup>.

4-{{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid was prepared according to the general method as outlined in example 1 (step7), starting from 4-{{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid ethylester (3.0 g, 5.15 mmol); 2.2 g of white powder. mp: 172 °C; Yield 77%; MS: 555 (M+H)<sup>+</sup>.

Starting from 4-{{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid (5.0 g, 9.0 mmol), and following the procedure as outlined in Example 1 (step 8), 1.8 g of 4-{{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid hydroxyamide was isolated as an yellow spongy solid. The dihydrochloride salt was prepared by dissolving the free amine with methanolic hydrochloric acid. mp: 124°C; Yield: 1.8 g (32%); MS: 570 (M+H)<sup>+</sup>.

#### Example 27

**4-{{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-(3-pentanyl)-piperidine-4-carboxylic acid hydroxyamide.**

4-{{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-(3-pentanyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 1 (step 6), starting from ethyl{{[4-(2-butynyloxy)phenyl]sulfonyl}acetate (8.8



g, 30.0 mmol) and bis-(2-chloro-ethyl)-(3-pentanyl)-amine dihydrochloride (7.4 g, 30 mmol), 3.5 g of product (brown oil) was isolated.. Yield 26%; MS: 436 (M+H)<sup>+</sup>

4-{{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-(3-pentanyl)-piperidine-4-carboxylic  
5 acid was prepared according to the general method as outlined in example 1 (step7),  
starting from 4-{{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-(3-pentanyl)-piperidine-4-  
carboxylic acid ethyl ester (3.0 g, 6.8 mmol); 2.5 g of spongy yellow solid. mp: 98 °C;  
Yield 90%; MS: 408 (M+H)<sup>+</sup>

10 Starting from 4-{{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-(3-pentanyl)-piperidine-4-  
carboxylic acid (2.5 g, 6.1 mmol), and following the procedure as outlined in Example 1  
(step 8), 1.8 g of 4-{{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-(3-pentanyl)-piperidine-4-  
carboxylic acid hydroxyamide was isolated as an yellow spongy solid. The hydrochloride  
salt was prepared by dissolving the free amine with methanolic hydrochloric acid. mp: 101-  
15 103°C; Yield: 1.1 g (42%); MS: 460 (M+H)<sup>+</sup>. <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 1.8 (t,  
6H), 1.5-1.7 (m, 6H), 1.9 (s, 3H), 2.3- 2.7 (m, 8H), 3.0 (m, 2H), 3.4 (s, 3H), 3.6 (d, 2H),  
4.9 (s, 2H), 7.21 (d, 2H), 7.8 (d, 2H), 9.3 (s, 1H), 9.8 (s, 1H), 11.2 (s, 1H).

#### Example 28

20 **1-(4-Methoxy-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-  
carboxylic acid hydroxyamide.**

1-(4-Methoxy-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-  
carboxylic acid ethyl ester was prepared according to the general method as outlined in  
example 1 (step 6), starting from (4-prop-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl  
25 ester (prepared as described in example 11, step 1 and 2)(10.0 g, 35.0 mmol) and 4-  
methoxy-benzyl)-bis-(2-chloro-ethyl)-amine hydrochloride (10.5 g, 35 mmol), 6.0 g  
of product (brown oil) was isolated.. Yield 36%; MS: 472 (M+H)<sup>+</sup>

1-(4-Methoxy-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-  
30 carboxylic acid was prepared according to the general method as outlined in example 1  
(step7), starting from 1-(4-Methoxy-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-

-72-

piperidine-4-carboxylic acid ethyl ester (6.0 g, 12.73 mmol); 5.0 g of spongy yellow solid. mp: 208 °C; Yield 92%; MS: 444 (M+H)<sup>+</sup>

Starting from 1-(4-Methoxy-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid (6.0 g, 13.5 mmol), and following the procedure as outlined in Example 1 (step 8), 2.0 g of 1-(4-Methoxy-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as an yellow spongy solid. The hydrochloride salt was prepared by dissolving the free amine with methanolic hydrochloric acid. mp: 150°C; Yield: 2.0 g (29%); MS: 459 (M+H)<sup>+</sup>. <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 2.3-2.8 (m, 6H), 3.3 (d, 2H), 3.5 (s, 3H), 4.2 (s, 2H), 5.0 (s, 2H), 7.3 (d, 2H), 7.5 (d, 2H), 7.6 (d, 2H), 7.7 (d, 2H), 10.9 (s, 1H), 11.2 (s, 1H).

#### Example 29

1-(4-Chloro-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide.

1-(4-Chloro-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester was prepared according to the general method as outlined in example 1 (step 6), starting from (4-prop-2-ynyloxy-benzenesulfonyl)-acetic acid ethyl ester (prepared as described in example 11, step 1 and 2)(10.0 g, 35.0 mmol) and 4-chloro-benzyl-bis-(2-chloro-ethyl)-amine hydrochloride (10.5 g, 35 mmol), 8.0 g of product (brown oil) was isolated.. Yield 48%; MS: 475 (M+H)<sup>+</sup>

1-(4-Chloro-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid was prepared according to the general method as outlined in example 1 (step7), starting from 1-(4-chloro-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid ethyl ester (6.0 g, 12.63 mmol); 5.0 g of spongy yellow solid. mp: 205 °C; Yield 92%; MS: 448 (M+H)<sup>+</sup>

Starting from 1-(4-Chloro-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid (6.0 g, 13.4 mmol), and following the procedure as

-73-

outlined in Example 1 (step 8), 2.0 g of 1-(4-chloro-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide was isolated as a yellow spongy solid. The hydrochloride salt was prepared by dissolving the free amine with methanolic hydrochloric acid. mp: 146°C; Yield: 4.0 g (59%); MS: 499 (M+H)<sup>+</sup>. <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 2.0-2.5 (m, 6H), 3.2 (d, 2H), 4.18 (s, 2H), 4.9 (s, 2H), 7.42 (d, 2H), 7.61 (d, 2H), 7.71 (d, 2H), 7.85 (d, 2H), 11.0 (s, 1H), 11.2 (s, 1H).

### Example 30

#### tert-butyl-4-([4-(2-butynyloxy)phenyl]sulfanyl)methyl)-4- [(hydroxyamino)carbonyl]-1-piperidinecarboxylate

10 Step 1: Piperidine-1,4-dicarboxylic acid tert-butyl ester ethyl ester

To a solution of ethyl isonipecotate (4.72g, 0.03 mmol) in 30mL of THF was added slowly di-tert-butyl dicarbonate (7.2g, 0.03 mmol) at room temperature. The resulting mixture was stirred for two hours and diluted with EtOAc. The organics  
15 were washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo. The residue was chromatographed on silica gel eluting with ethyl acetate:hexanes (1:9) to provide 7.52g (97%) of the desired product as a colorless oil. Electrospray Mass Spec : 258.3 (M+H)<sup>+</sup>

20 Step 2: 1-(tert-Butyl) 4-ethyl 4-(iodomethyl)piperidine-1,4-dicarboxylate

To a solution of piperidine-1,4-dicarboxylic acid tert-butyl ester ethyl ester (12.8 g, 49.74 mmol) in 73mL of dry THF under N<sub>2</sub> atmosphere at -42°C was added 24.87mL (49.74mmol) of 2M Lithium diisopropylamine in heptane/THF/ethylbenzene dropwise to not exceed -40°C. After one hour, 4.0mL (49.74mmol) of diiodomethane was  
25 added and the solution was warmed to ambient temperature overnight. The resulting solution was diluted with H<sub>2</sub>O and extracted with ethyl acetate. The organics were washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo to provide 18.84g(95%) of the desired product as a brown oil. Electrospray Mass Spec : 398.2 (M+H)<sup>+</sup>

30

-74-

## Step 3: 4-But-2-ynyloxy-benzenesulfonic acid sodium salt:

To a solution of 4-hydroxybenzenesulfonate sodium salt (52.35g, 0.225) in 1L of isopropanol and 225mL of a 1.0N solution of sodium hydroxide was added 59.96g (0.45mol) of 1 bromo-2-butyne. The resulting mixture was heated to 70° for 15h and then the isopropanol was removed by evaporation in vacuo. The resulting white precipitate was collected by filtration, washed with isopropanol and ether and dried in vacuo to give 45.08g (81%) of the desired product as a white solid.

## Step 4: 4-But-2-ynyloxy-benzenesulfonyl chloride

To a stirred solution of oxalyl chloride (47.8 ml, 0.545 mol) at 0°C in 240mL of CH<sub>2</sub>Cl<sub>2</sub> was added a DMF (43.0 ml) solution of 4-but-2-ynyloxy-benzenesulfonic acid sodium salt in a drop wise manner. The reaction mixture was stirred at 0 °C for 30min and then let warm to room temperature and stirred for 18h. The reaction was then poured into ice and extracted with hexanes. The organics were washed with water and brine, dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo to provide 42.0g (95%) of the desired product as yellow solid.

## Step 5: 4-But-2-ynyloxy-benzenethiol

To a solution of 11.8g (0.045mol) of triphenylphosphine in 10mL of CH<sub>2</sub>Cl<sub>2</sub> and 0.3mL of DMF was added dropwise a solution of 4-but-2-ynyloxy-benzenesulfonyl chloride in 15 mL CH<sub>2</sub>Cl<sub>2</sub>. Stirred at room temperature for two hours, added 5mL of 1N HCl, stirred for 30 min., and then added 15mL of brine. The organics were separated and concentrated in vacuo. The residue was diluted with ether and filtered the insolubles. The filtrate was washed with 2.5N NaOH and the aqueous solution separated, acidified and extracted with ether. The organics were washed with H<sub>2</sub>O, brine, dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo to give 1.54g (58%) of the desired product as a pale yellow oil.

-75-

Step 6: 1-(tert-butyl) 4-ethyl 4-([4-(2-butynyloxy)phenyl]sulfanyl) methyl)-1,4-piperidinedicarboxylate

A mixture of 0.294g (0.74mmol) of 1-(tert-Butyl) 4-ethyl 4-(iodomethyl)-piperidine-1,4-dicarboxylate, 0.145 (0.814mmol) of 4-but-2-ynyloxy-benzenethiol and  
5 0.204g (1.48mmol) of  $K_2CO_3$  in 2.0mL of DMF was stirred at room temperature for 18 h. The resulting mixture was diluted with EtOAc, washed with  $H_2O$ , brine, dried over  $MgSO_4$  and concentrated in vacuo. The residue was chromatographed on silica gel eluting with EtOAc:Hexanes (1:19) to provide 0.328g (99%) of the desired product as a colorless oil. Electrospray Mass Spec : 448.3 (M+H)<sup>+</sup>

10

Step 7: 4-(4-But-2-ynyloxy-phenylsulfanylmethyl)-piperidine-1,4- dicarboxylic acid mono-tert-butyl ester

A mixture of 0.288g (0.0643mmol) of 1-(tert-butyl) 4-ethyl 4-([4-(2-butynyloxy)phenyl]sulfanyl) methyl)-1,4-piperidinedicarboxylate, 3.25mL of 1N NaOH  
15 3.25mL of THF and 3.25mL of MeOH was heated to reflux for 3h. The organics were removed and the residue was diluted with  $H_2O$ , acidified and extracted with EtOAc. The organics were washed with  $H_2O$ , brine, dried over  $mgSO_4$ , filtered and concentrated in vacuo to provide 0.241g (89%) of the desired product as an off white gum. Electrospray Mass Spec: 464.3 (M+FA-H)<sup>-</sup>

20 Step 8: WAY 173665 tert-butyl-4-([4-(2-butynyloxy)phenyl]sulfanyl)methyl)-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxylate

To a solution of 0.204g (0.49mmol) of 4-(4-But-2-ynyloxy-phenylsulfanylmethyl)-piperidine-1,4- dicarboxylic acid mono-tert-butyl ester, 0.079g (0.58mmol) of 1-hydroxybenzotriazole in 2.5mL of DMF was added 0.112g (0.84mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, and stirred at room  
25 temperature for 1h. Then added 0.3mL of 50% aqueous hydroxylamine and stirred for 18h. The resulting mixture was diluted with EtOAc, washed with  $H_2O$ , brine, dried over  $MgSO_4$ , filtered and concentrated in vacuo. The residue was chromatographed on silica gel eluting with 1.5% MeOH/  $CH_2Cl_2$  to provide 0.077g (36%) of the desired  
30 product as a white solid. Electrospray Mass Spec: 435.2 (M+H)<sup>+</sup>

**Example 31****4-([4-(But-2-ynyloxy)phenyl]thio)methyl)-N-hydroxypiperidine-4-carboxamide**

To a solution of 0.143g (0.033mmol) of tert-butyl-4-([4-(2-butyloxy)phenyl]sulfonyl)methyl)-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxylate in 5mL of  $\text{CH}_2\text{Cl}_2$  and 1mL of MeOH was added 5mL of 4M HCl in dioxane and stirred for 1h. The reaction was concentrated in vacuo and the residue was triturated with ether and filtered to provide 0.093g (76%) of the desired product as a pale orange solid. Electrospray Mass Spec: 335.3 (M+H)<sup>+</sup>

10

**Example 32****tert-Butyl-4-([4-(2-butyloxy)phenyl]sulfonyl)methyl)-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxylate**

To a slurry of tert-butyl-4-([4-(2-butyloxy)phenyl]sulfonyl)methyl)-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxylate (0.24g, 0.55 mmol) at 0°C in 7mL of MeOH was added dropwise 3.5mL of 30% hydrogen peroxide. The reaction was allowed to warm to room temperature and stirred for 18h. The reaction was cooled to 0°C and quenched with 3.5mL of a saturated solution of  $\text{Na}_2\text{SO}_3$ . The organics were removed and the aqueous solution was extracted with  $\text{CH}_2\text{Cl}_2$ . The organics were washed with  $\text{H}_2\text{O}$ , brine, dried over  $\text{MgSO}_4$ , filtered and concentrated in vacuo. The residue was triturated with ether to provide 0.166g (67%) of the desired product as an off white solid. Electrospray Mass Spec: 451.3 (M+H)<sup>+</sup>

15

20

**Example 33****4-([4-(2-Butyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-piperidinecarboxamide**

25

4-([4-(2-Butyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-piperidinecarboxamide was prepared according to the general method as outlined in Example 31. Starting from tert-butyl-4-([4-(2-butyloxy)phenyl]sulfonyl)methyl)-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxylate (0.082 g, 0.18 mmol), 0.066g (95%) of the desired product was isolated as a white solid. Electrospray Mass Spec: 351.2 (M+H)<sup>+</sup>

30

-77-

**Example 34****tert-Butyl-4-([4-(but-2-ynyloxy)phenyl]sulfonyl)methyl)-4-  
[(hydroxyamino)carbonyl]piperidine-1-carboxylate**

To a solution of tert-butyl-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-  
5 [(hydroxyamino)carbonyl]-1-piperidinecarboxylate (0.422g, 0.97 mmol) in 8mL of  
MeOH, 4mL of CH<sub>2</sub>Cl<sub>2</sub> and 2mL of THF was added a solution of 1.79g (2.91mmol)  
of OXONE in 8mL of H<sub>2</sub>O and stirred at room temperature for 18h. The solid was  
filtered and the filtrate was concentrated in vacuo. The residue was diluted with  
EtOAc, washed with H<sub>2</sub>O, brine, dried over MgSO<sub>4</sub>, filtered, and concentrated to  
10 provide 0.351g (77%) of the desired product as a white solid. Electrospray Mass Spec:  
467.3 (M+H)<sup>+</sup>

**Example 35****tert-butyl-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-  
15 [(hydroxyamino)carbonyl]-1-piperidinecarboxylate**

tert-Butyl-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-[(hydroxyamino)-  
carbonyl]-1-piperidinecarboxylate was prepared according to the general method as  
outlined in Example 31. Starting from tert-Butyl-4-([4-(but-2-ynyloxy)-  
phenyl]sulfonyl)methyl)-4-[(hydroxyamino)carbonyl]piperidine-1-carboxylate (0.10g,  
20 0.214mmol) 0.074g (86%) of the desired product was isolated as white solid.  
Electrospray Mass Spec: 367.3 (M+H)<sup>+</sup>

**Example 36****1-Acetyl-4-[[[4-(2-butynyloxy)phenyl]sulfonyl)methyl]-N-hydroxy-4-  
25 piperidinecarboxamide**

Step 1: 4-[[[4-(2-Butynyloxy)phenyl]sulfonyl)methyl]-1,4- piperidine dicarboxylic  
acid, 1-tert-butyl 4-ethyl ester

To a solution of 1-(tert-butyl) 4-ethyl 4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-  
1,4- piperidinedicarboxylate (1.66 g, 3.7 mmol) (prepared in example 30, step 6) in  
30 20mL of CH<sub>2</sub>Cl<sub>2</sub> was added tetrabutylammonium oxone (17.38g, 14.7 mmol) and  
stirred at room temperature for 18h. The reaction was concentrated in vacuo and the

-78-

residue was diluted with EtOAc, washed with H<sub>2</sub>O, 5% KHSO<sub>4</sub>, brine, dried over MgSO<sub>4</sub>, filtered and concentrated to provide 1.69g (95%) of the desired product as a pale yellow gum. Electrospray Mass Spec: 480.3 (M+H)<sup>+</sup>

- 5 Step 2 4-[[[4-(2-Butynyloxy)phenyl]sulfonyl]methyl]-4-piperidinecarboxylic acid, ethyl ester

4-[[[4-(2-Butynyloxy)phenyl]sulfonyl]methyl]-4-piperidinecarboxylic acid ethyl ester was prepared according to the general method as outlined in Example 31. Starting from 4-[[[4-(2-Butynyloxy)phenyl]sulfonyl]methyl]-1,4-piperidinedicarboxylic acid 1-tert-butyl 4-ethyl ester (1.62g 3.4mmol), 1.335g (95%) of the desired product  
10 was isolated as a tan solid. Electrospray Mass Spec: 380.2 (M+H)<sup>+</sup>

Step 3: 1-Acetyl-4-[[[4-(2-butynyloxy)phenyl]sulfonyl]methyl]-4-piperidinecarboxylic acid, ethyl ester

- 15 To a solution of 4-[[[4-(2-Butynyloxy)phenyl]sulfonyl]methyl]-4-piperidine-carboxylic acid ethyl ester (0.24g, 0.576mmol), triethylamine (0.32 ml) and catalytic amount of 4-Dimethylaminopyridine in 6.0mL of CH<sub>2</sub>Cl<sub>2</sub> was added a solution of acetyl chloride (0.068 ml, 0.864 mmol) in 1.0mL of CH<sub>2</sub>Cl<sub>2</sub>. The reaction stirred at room temperature for 4h and washed with H<sub>2</sub>O, brine, dried over MgSO<sub>4</sub>, filtered  
20 through a pad of silica gel and concentrated to provide 0.242g (100%) of the desired product as a colorless gum. Electrospray Mass Spec: 422.2 (M+H)<sup>+</sup>

Step 4: 1-Acetyl-4-(4-but-2-ynyloxy-benzenesulfonylmethyl)-piperidine-4- carboxylic acid

- 25 1-Acetyl-4-(4-but-2-ynyloxy-benzenesulfonylmethyl)-piperidine-4- carboxylic acid was prepared according to the general method as outlined in Example 30, (step 7). Starting from 1-Acetyl-4-[[[4-(2-butynyloxy)phenyl]sulfonyl]methyl]-4-piperidine-carboxylic acid, ethyl ester (0.22 g, 0.524 mmol), 0.141 g of the desired product was isolated as a pale yellow solid. Electrospray Mass Spec: 438.2 (M+FA-H)<sup>-</sup>

30



-79-

Step 5: 1-Acetyl-4-[[[4-(2-butynyloxy)phenyl]sulfonyl]methyl]-N-hydroxy-4-piperidinecarboxamide

1-Acetyl-4-[[[4-(2-butynyloxy)phenyl]sulfonyl]methyl]-N-hydroxy-4-piperidinecarboxamide was prepared according to the general method as outlined in  
5 Example 30 (step 8). Starting from 1-Acetyl-4-(4-but-2-ynyloxy-benzenesulfonylmethyl)-piperidine-4-carboxylic acid, (0.122 g, 0.31 mmol) 0.048g (38%) of the desired product was isolated as a pale yellow solid. Electrospray Mass Spec: 409.2 (M+H)<sup>+</sup>

10

### Example 37

#### 1-(2-Butynyl)-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-piperidinecarboxamide hydrochloride

Step 1: 1-(2-Butynyl)-4-[[[4-(2-butynyloxy)phenyl]sulfonyl]methyl]-4-piperidinecarboxylic acid, ethyl ester

15 A mixture of 4-[[[4-(2-Butynyloxy)phenyl]sulfonyl]methyl]-4-piperidinecarboxylic acid ethyl ester (0.208 g, 0.5 mmol), 1-bromo-2-butyne (0.044 ml, 0.53 mmol) and K<sub>2</sub>CO<sub>3</sub> (0.138 g, 1.0 mmol) in 5.0 mL of DMF was stirred at room temperature for 6h. The reaction was diluted with EtOAc and washed with H<sub>2</sub>O, brine, dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo. The residue was  
20 chromatographed on silica gel eluting with EtOAc:hexanes (1:1) to provide 0.183g (85%) of the desired product as a pale yellow gum. Electrospray Mass Spec: 432.2 (M+H)<sup>+</sup>

25 Step 2: 1-(2-Butynyl)-4-[4-(2-butynyloxy)benzenesulfonylmethyl]-piperidine-4-carboxylic acid

1-(2-Butynyl)-4-[4-(2-butynyloxy)benzenesulfonylmethyl]-piperidine-4-carboxylic acid was prepared according to the general method as outlined in example  
30 (step 7). Starting from 1-(2-Butynyl)-4-[[[4-(2-butynyloxy)phenyl]sulfonyl]methyl]-4-piperidinecarboxylic acid, ethyl ester, (0.153 g, 0.354 mmol), 0.12 g (84%) of the desired product was isolated as a white solid. Electrospray Mass Spec: 404.2 (M+H)<sup>+</sup>

Step 3: 1-(2-Butynyl)-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-piperidinecarboxamide hydrochloride

1-(2-Butynyl)-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-piperidinecarboxamide hydrochloride was prepared according to the general method as outlined in Example 30 (step 8). Starting from 1-(2-butynyl)-4-[4-(2-butynyloxy)-benzenesulfonylmethyl]-piperidine-4-carboxylic acid, (0.15g, 0.34mmol), 0.05g of the desired product, which was dissolved in 1.0mL of CH<sub>2</sub>Cl<sub>2</sub> and treated with 0.225mL of 1M HCl in CH<sub>2</sub>Cl<sub>2</sub>. The solution was stirred for 1h, and concentrated in vacuo. The residue was triturated with ether to provide 0.044g (28%) of the hydrochloride of the desired product as a beige solid. Electrospray Mass Spec: 419.2 (M+H)<sup>+</sup>

#### Example 38

N-1-(tert-Butyl)-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-4-hydroxy-1,4-[4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-4-hydroxy-1,4-l[sulfonyl)methyl)-N-4-hydroxy-1,4-piperidinedicarboxamide

Step 1: 1-tert-Butylcarbamoyl-4-(4-but-2-ynyloxy-benzenesulfonylmethyl)-piperidine-4-carboxylic acid ethyl ester

To a solution of tert-butyliisocyanate (0.097 ml, 0.85 mmol) in 8.0mL of CH<sub>2</sub>Cl<sub>2</sub>, was added 4-[[[4-(2-Butynyloxy)phenyl]sulfonyl)methyl]-4-piperidine-carboxylic acid ethyl ester (prepared from example 36, step 2) (0.337 g, 0.81 mmol) and triethylamine (0.135 ml, 0.97 mmol) and stirred at room temperature for 2h. The reaction was diluted with CH<sub>2</sub>Cl<sub>2</sub> and washed with H<sub>2</sub>O, brine, dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo. The residue was triturated with ether:hexanes (1:1) to provide 0.284g (73%) of the desired product as a white solid. Electrospray Mass Spec: 479.2 (M+H)<sup>+</sup>

Step 2: 1-[(tert-Butylamino)carbonyl]-4-([4-(2-butynyloxy)phenyl] sulfonyl)methyl)-4-onyl]-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-piperidinecarboxylic acid

1-[(tert-Butylamino)carbonyl]-4-([4-(2-butynyloxy)phenyl]sulfonyl)-methyl)-4-onyl]-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-piperidine

-81-

carboxylic acid was prepared according to the general method as outlined in Example 30 (step 7). Starting from 1-tert-butylcarbonyl-4-(4-but-2-ynyloxy-benzene-sulfonylmethyl)-piperidine-4-carboxylic acid ethyl ester (0.259g, 0.54 mmol), 0.169 g, (69%) of the desired product was isolated as white solid. Electrospray Mass Spec: 451.4 (M+H)<sup>+</sup>

Step 3: N-1-(tert-Butyl)-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-4-hydroxy-1,4-[4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-4-hydroxy-1,4-l[sulfonyl)methyl)-N-4-hydroxy-1,4-piperidinedicarboxamide

N-1-(tert-Butyl)-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-4-hydroxy-1,4-[4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-4-hydroxy-1,4-l[sulfonyl)methyl)-N-4-hydroxy-1,4-piperidinedicarboxamide was prepared according to the general method as outlined in Example 30 (step 8). Starting from 1-[(tert-Butylamino)carbonyl]-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-onyl]-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-piperidinecarboxylic acid (0.149g, 0.33 mmol), 0.077 g of the desired product was isolated as pale yellow solid. Electrospray Mass Spec: 466.3 (M+H)<sup>+</sup>

### Example 39

Methyl 4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxylate

Step 1: 4-(4-But-2-ynyloxy-benzenesulfonylmethyl)-piperidine-1,4-dicarboxylic acid ethyl ester methyl ester

To a solution of 4-[[[4-(2-Butynyloxy)phenyl]sulfonyl]methyl]-4-piperidine-carboxylic acid ethyl ester (0.354 g, 0.85 mmol) in 1.0mL of CH<sub>2</sub>Cl<sub>2</sub> under N<sub>2</sub> atmosphere was added dropwise a solution of N,O-bis(trimethylsilyl)acetamide (0.462 ml, 1.87 mmol) in 0.5mL of CH<sub>2</sub>Cl<sub>2</sub> and stirred for 1h. The reaction was cooled to 0°C and added dropwise a solution of 0.079mL (1.02mmol) of methylchloroformate in 0.5mL of CH<sub>2</sub>Cl<sub>2</sub>. The reaction was allowed to stir at room temperature for 1h and cooled to 0°C, quenched with pH7 buffer solution and extracted with EtOAc. The organics was washed with H<sub>2</sub>O, brine, dried over MgSO<sub>4</sub>, filtered, and concentrated in

-82-

vacuo. The residue was chromatographed on silica gel eluting with EtOAc:hexanes (1:2) to provide 0.315g (85%) of the desired product as a colorless oil. Electrospray Mass Spec: 438.3 (M+H)<sup>+</sup>

- 5 Step 2: 4-(4-But-2-ynyloxy-benzenesulfonylmethyl)-piperidine-1,4- dicarboxylic acid monomethyl ester

4-(4-But-2-ynyloxy-benzenesulfonylmethyl)-piperidine-1,4-dicarboxylic acid monomethyl ester was prepared according to the general method as outlined in Example 30 (step 7). Starting from 4-(4-But-2-ynyloxy-benzenesulfonylmethyl)-  
10 piperidine-1,4- dicarboxylic acid ethyl ester methyl ester (0.277 g, 0.633 mmol), 0.175g (67%) of the desired product was isolated as white solid. Electrospray Mass Spec: 410.2 (M+H)<sup>+</sup>

- Step 3: Methyl 4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-4-[(hydroxyamino)-  
15 carbonyl]-1-piperidinecarboxylate

Methyl-4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-4-[(hydroxyamino)-carbonyl]-1-piperidinecarboxylate was prepared according to the general method as outlined in Example 30 (step 8). Starting from 4-(4-But-2-ynyloxy-benzenesulfonylmethyl)-piperidine-1,4-dicarboxylic acid monomethyl ester (0.15 g, 0.366 mmol),  
20 0.053g (34%) of the desired product was isolated as a white solid. Electrospray Mass Spec: 425.3 (M+H)<sup>+</sup>

#### Example 40

- 25 **Benzyl 4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxylate**

Step 1: 4-(4-But-2-ynyloxy-benzenesulfonylmethyl)-piperidine-1,4-dicarboxylic acid benzyl ester ethyl ester

4-(4-But-2-ynyloxy-benzenesulfonylmethyl)-piperidine-1,4- dicarboxylic acid benzyl ester ethyl ester was prepared according to the general method as outlined in  
30 Example 39 (step 1). Starting from 4-[[[4-(2-Butynyloxy)phenyl]sulfonyl]methyl]-4-

-83-

piperidinecarboxylic acid ethyl ester (0.312 g, 0.75), 0.337g (87%) of the desired product was isolated as colorless oil. Electrospray Mass Spec: 514.2 (M+H)<sup>+</sup>

Step 2: 4-(4-But-2-ynyloxy-benzenesulfonylmethyl)-piperidine-1,4- dicarboxylic acid  
5 monobenzyl ester

4-(4-But-2-ynyloxy-benzenesulfonylmethyl)-piperidine-1,4- dicarboxylic acid  
monobenzyl ester was prepared according to the general method as outlined in  
Example 30 (step 7). Starting from 4-(4-But-2-ynyloxy-benzenesulfonylmethyl)-  
piperidine-1,4- dicarboxylic acid benzyl ester ethyl ester (0.32g, 0.623 mmol) , 0.2 g of  
10 the desired product was isolated as white solid. Electrospray Mass Spec: 484.2  
(M-H)<sup>-</sup>

Step 3: Benzyl 4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-[(hydroxyamino)-  
carbonyl]-1-piperidinecarboxylate

15 Benzyl 4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-[(hydroxyamino)-  
carbonyl]-1-piperidinecarboxylate was prepared according to the general method as  
outlined in Example 30, (step 8). Starting from 4-(4-But-2-ynyloxy-benzene  
sulfonylmethyl)-piperidine-1,4-dicarboxylic acid monobenzyl ester (0.18 g, 0.37  
mmol), 0.106 g (57%) of the desired product was isolated as off-white solid.  
20 Electrospray Mass Spec: 501.3 (M+H)<sup>+</sup>

#### Example 41

**1-Benzyl-4-([4-(2-butynyloxy)phenyl] sulfonyl) methyl)-N-hydroxy-4-  
butynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-piperidinecarboxamide**

25 Step 1: Ethyl-1-benzyl-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-4-(2-butynyl-  
oxy)phenyl]sulfonyl)methyl)-4-piperidinecarboxylate  
Ethyl-1-benzyl-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-4-(2-butynyloxy)-  
phenyl]sulfonyl)methyl)-4-piperidinecarboxylate was prepared according to the  
general method as outlined in Example 37 (step 1). Starting from 4-[[[4-(2-Butynyl-  
30 oxy)phenyl]sulfonyl]methyl)-4-piperidinecarboxylic acid ethyl ester (prepared in

Example 36, step 2) (0.312g, 0.75 mmol), 0.265g of the desired product was isolated as white solid. Electrospray Mass Spec: 470.2 (M+H)<sup>+</sup>

Step 2: 1-Benzyl-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-piperidine-carboxylic-benzyl-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-piperidine-carboxylic acid

1-Benzyl-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-piperidinecarboxylic acid was prepared according to the general method as outlined in Example 30 (step 7). Starting from Ethyl-1-benzyl-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-piperidinecarboxylate (0.25g, 0.53 mmol), 0.227g (90%) of the desired product was isolated as a white solid. Electrospray Mass Spec: 442.2 (M+H)<sup>+</sup>

Step 3: 1-Benzyl-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-utynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-piperidinecarboxamide 1-Benzyl-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-butynyl-oxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-piperidinecarboxamide was prepared according to the general method as outlined in Example 30 (step 8). Starting from 1-Benzyl-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-piperidinecarboxylic-enzyl-4-([4-(2 butynyloxy)phenyl]sulfonyl)methyl)-4-piperidinecarboxylic acid (0.211g, 0.44 mmol), 0.108g of the desired product was isolated as white solid.. Electrospray Mass Spec: 457.2 (M+H)<sup>+</sup>

25

#### Example 42

4-([4-(2-Butynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxamide

Step 1: Ethyl 4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-1-[(2,2,5- trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxylate

Ethyl 4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-1-[(2,2,5- trimethyl-1,3-dioxan 5-yl)carbonyl]-4-piperidinecarboxylate was prepared according to the general method

-85-

as outlined in Example 30 (step 8). Starting from 4-[[[4-(2-Butynyloxy)-phenyl]sulfonyl]methyl]-4-piperidinecarboxylic acid ethyl ester (0.333g, 0.8 mmol) and 2,2,5-trimethyl-(1,3)dioxane-5-carboxylic acid (0.168g, 0.96 mmol), 0.339g (79%) of the desired product was isolated as a white solid. Electrospray Mass Spec:  
5 536.1 (M+H)<sup>+</sup>

Step 2: 4-([4-(2-Butynyloxy)phenyl]sulfonyl)methyl-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxylic acid  
4-([4-(2-Butynyloxy)phenyl]sulfonyl)methyl-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)-  
10 carbonyl]-4-piperidinecarboxylic acid was prepared according to the general method as outlined in Example 30 (step 7). Starting from ethyl 4-([4-(2-butynyloxy)-phenyl]sulfonyl)methyl-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidine-carboxylate (0.299g, 0.558 mmol), 0.235g (83%) of the desired product was isolated as white solid. Electrospray Mass Spec: 506.2 (M-H)<sup>-</sup>  
15

Step 3: 4-([4-(2-Butynyloxy)phenyl]sulfonyl)methyl-N-hydroxy-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxamide  
4-([4-(2-Butynyloxy)phenyl]sulfonyl)methyl-N-hydroxy-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxamide was prepared according  
20 to the general method as outlined in Example 30 (step 8). Starting from 4-([4-(2-Butynyloxy)phenyl]sulfonyl)methyl-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxylic acid (0.22g, 0.433 mmol), 0.16 g of the desired product was isolated as white solid. Electrospray Mass Spec: 523.2 (M+H)<sup>+</sup>

25

### Example 43

4-([4-(2-Butynyloxy)phenyl]sulfonyl)methyl-N-hydroxy-1-[3-hydroxy-2-(hydroxymethyl)-2-methylpropanoyl]-4-piperidinecarboxamide

A mixture of 4-([4-(2-Butynyloxy)phenyl]sulfonyl)methyl-N-hydroxy-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxamide (0.106 g, 0.2  
30 mmol) and 2mL of 1N HCl in 2mL of THF was stirred at room temperature for 4h. The reaction was diluted with EtOAc, washed with H<sub>2</sub>O, saturated NaHCO<sub>3</sub>, brine,

-86-

dried over  $\text{MgSO}_4$ , filtered, and concentrated in vacuo. The residue was triturated with ether to provide 0.67g (71%) of the desired product as an off white solid. Electrospray Mass Spec: 483.2 (M+H)<sup>+</sup>

5

**Example 44**

**1-[Amino(imino)methyl]-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-l]-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-oxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-piperidinecarboxamide**

Step 1: N,N'-t-Boc-protected thiourea: To a stirred solution of thiourea (0.57g, 7.5 mmol) in 150mL of THF under  $\text{N}_2$  at 0°C was added 60% NaH (1.35g, 33.8 mmol) in mineral oil. After 5 minutes, the ice bath was removed and the reaction mixture was allowed to stir at room temperature for 10 minutes. The reaction mixture was cooled to 0°C and 3.6g (16.5mmol) of di-tert-butyl dicarbonate was added. After 30 minutes, the ice bath was removed and the reaction was stirred for 2h. The reaction was then quenched with saturated  $\text{NaHCO}_3$  solution, poured into water and extracted with 3x EtOAc. The organics were washed with  $\text{H}_2\text{O}$ , brine, dried over  $\text{MgSO}_4$ , filtered, and concentrated in vacuo. The residue was triturated with hexane to provide 1.72g (83%) of the desired product as a white solid.

Step 2: tert-Butyl 4-[(tert-butoxyamino)carbonyl]-4-([4-(2-yloxy)phenyl]sulfonyl)methyl)-1-piperidinecarboxylate

tert-Butyl 4-[(tert-butoxyamino)carbonyl]-4-([4-(2-yloxy)phenyl]sulfonyl)-methyl)-1-piperidinecarboxylate was prepared according to the general method as outlined in Example 30 (step 8). Starting from 4-(4-but-2-ynyloxy-benzenesulfonyl methyl)-piperidine-1,4- dicarboxylic acid mono-tert-butyl ester (2.53 g, 5.6 mmol) and O-tert-butyl-hydroxylamine hydrochloride (1.4 g, 11.2 mmol), 2.31 g (79%) of the desired product was isolated as a white solid. Electrospray Mass Spec: 523.2 (M+H)<sup>+</sup>



-87-

Step 3: N-(tert-Butoxy)-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-[4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-piperidinecarboxamide

To a solution of tert-Butyl 4-[(tert-butoxyamino)carbonyl]-4-([4-(2-yloxy)-phenyl]sulfonyl)methyl)-1-piperidinecarboxylate (3.0 g, 5.5 mmol) in 6mL of CH<sub>2</sub>Cl<sub>2</sub> was added of trimethylsilyltrifluoromethylsulfonate (1.1 mL, 6.05 mmol) followed by 0.7mL of 2,6-lutidine. The reaction was stirred for 1h and diluted with CH<sub>2</sub>Cl<sub>2</sub>. The organics were washed with H<sub>2</sub>O, saturated NaHCO<sub>3</sub>, brine, dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide 2.01g (86%) of the desired product as an off white solid. Electrospray Mass Spec: 423.2 (M+H)<sup>+</sup>

10

Step 4: [[4-[(tert-Butoxyamino)carbonyl]-4-[[[4-(2-[4-(2-butynyloxy)phenyl]-sulfonyl)methyl)-4-piperidinecarboxamide butoxycarbonyl]amino]methylene]carbamic acid, tert-butyl ester

To a mixture of N-(tert-Butoxy)-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-[4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-piperidinecarboxamide (0.127g, 0.3mmol), the di-t-boc-protected thiourea (obtained from step 1) (0.091g, 0.33 mmol) and triethylamine (0.092 mL) in 3mL of DMF was added mercury(II) chloride (0.09g, 0.33 mmol) and stirred for 1h at 0°C. The reaction was diluted with EtOAc and filtered through a pad of celite. The organics were washed with H<sub>2</sub>O, brine, dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo. The residue was triturated with hexanes to provide the desired product as a white solid. . Electrospray Mass Spec: 665.5 (M+H)<sup>+</sup>

20

Step 5: 1-[Amino(imino)methyl]-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-[4-(2-butynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-oxy)phenyl]sulfonyl)methyl)-N-hydroxy-4-piperidinecarboxamide

25

A mixture of [[4-[(tert-Butoxyamino)carbonyl]-4-[[[4-(2-[4-(2-butynyloxy)-phenyl]sulfonyl)methyl)-4-piperidinecarboxamidebutoxycarbonyl]amino]methylene]-carbamic acid, tert-butyl ester (0.135g, 0.2 mmol) and 3mL of trifluoroacetic acid in 2mL of CH<sub>2</sub>Cl<sub>2</sub> was heated at 60°C for 24h. The reaction was concentrated in vacuo and was prep HPLC to provide 0.032g (31%) of the desired product as a beige solid. Electrospray Mass Spec: 409.3 (M+H)<sup>+</sup>

30

## 5

10

15

20

30

-89-


ethyl-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-1-(4-hydroxy-2-butynyl)-4-piperidinecarboxylate (0.115g, 0.257 mmol), 0.08g (74%) of the desired product was isolated as white solid. Electrospray Mass Spec: 420.4 (M+H)<sup>+</sup>

- 5 Step 4: 4-([4-(2-Butynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(4-hydroxy-2-butynyl)-4-piperidinecarboxamide

4-([4-(2-Butynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(4-hydroxy-2-butynyl)-4-piperidinecarboxamide was prepared according to the general method as outlined in Example 30 (step 8). Starting from 4-([4-(2-Butynyloxy)phenyl]sulfonyl)methyl)-1-(4-hydroxy-2-butynyl)-4-piperidinecarboxylic acid (0.073g, 0.174 mmol), 0.026g (34%) of the desired product was isolated as white solid. Electrospray Mass Spec: 435.3 (M+H)<sup>+</sup>

15

Methods for the solution phase synthesis of the compounds of the present invention is as shown in the following scheme.



20

#### Example 46

**4-([4-(2-Butynyloxy)phenyl]sulfonyl)methyl)-1-ethyl-N-hydroxypiperidine-4-carboxamide trifluoroacetic acid salt**

- 25 Step A: A solution of N-(tert-butoxy)-4-([4-(2-butynyloxy)phenyl]sulfonyl)methyl)-4-piperidinecarboxamide (0.097g, 0.23mmol), ethyl iodide (0.019mL, 0.24mmol) and triethylamine (0.096mL, 0.69mmol) in 2mL of CH<sub>2</sub>Cl<sub>2</sub> was shaken at room temperature for 18h and then concentrated in vacuo.

-90-

Step B: A solution of the residue from Step A in 1mL of CH<sub>2</sub>Cl<sub>2</sub> and 1mL of trifluoroacetic acid was heated at 50°C for 2h and then concentrated in vacuo to provide the desired product.

5        The following hydroxamic acids were synthesized according to the procedures of 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-1-ethyl-N-hydroxypiperidine-4-carboxamide trifluoroacetic acid salt using the appropriate reagents.

Example 47: Reagent - 0.029mL (0.24mmol) of 2-chloro-5-(chloromethyl)thiophene-  
10 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-1-[(5-chlorothien-2-yl)methyl]-N-hydroxypiperidine-4-carboxamide trifluoroacetic acid salt

Example 48: Reagent - 0.0496g (0.24mmol) of 4-picolyl chloride hydrochloride  
15 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(pyridin-4-ylmethyl)-piperidine-4-carboxamide trifluoroacetic acid salt

#### Example 49

4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(pyridin-3-ylcarbonyl)piperidine-4-carboxamide trifluoroacetic acid salt

20        Step A: A solution of N-(tert-butoxy)-4-([4-92-butynyloxy)phenyl]sulfonyl)methyl)-4-[4-(2-butynyloxy)phenyl]sulfonyl)methyl-4-piperidine carboxamide (0.097g, 0.23mmol), triethylamine (0.064mL, 0.64mmol), nicotinoyl chloride hydrochloride (0.061g, 0.34mmol), and 4-dimethylaminopyridine (0.002 g) in 2mL of CH<sub>2</sub>Cl<sub>2</sub> was shaken at room temperature for 18h and then concentrated in vacuo.

25

Step B: same as Step B of Example 46.

The following hydroxamic acids were synthesized according to the procedures of 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(pyridin-3-ylcarbonyl)-piperidine-4-carboxamide trifluoroacetic acid salt using the appropriate reagents.

30

-91-

**Example 50****Reagent - 0.04mL (0.276mmol) of benzoyl chloride****1-Benzoyl-4-([4-(but-2-ynyloxy)phenyl]sulfonyl)methyl-N-hydroxypiperidine-4-carboxamide**

5

**Example 51****Reagent - 0.037mL (0.276mmol) of 2-thiophenecarbonyl chloride****4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl-N-hydroxy-1-(thien-2-ylcarbonyl) piperidine-4-carboxamide**

10

**Example 52****4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl-N-1-ethyl-N-4-hydroxypiperidine-1,4-dicarboxamide**

Step A: A solution of N-(tert-butoxy)-4-([4-92-butynyloxy)phenyl]sulfonyl)methyl)-

15 4-[4-(2-butynyloxy)phenyl]sulfonyl)methyl-4-piperidinecarboxamide (0.097g, 0.23mmol), triethylamine (0.064mL, 0.64mmol) and ethyl isocyanate (0.02mL, 0.253mmol) in 2mL of CH<sub>2</sub>Cl<sub>2</sub> was shaken at room temperature for 18h and then concentrated in vacuo.

Step B: same as Step B of Example 46.

20 The following hydroxamic acids were synthesized according to the procedures of Example 52.using the appropriate reagents.

**Example 53****Reagent - 0.275mL (0.253mmol) of phenylisocyanate**

25 **4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl-N-4-hydroxy-N-1-phenylpiperidine-1,4-dicarboxamide**

**Example 54****Reagent - 0.32mL (0.253mmol) of diethylcarbamyl chloride**

30 **4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl-N-1-,N-1-diethyl-N-4-hydroxypiperidine-1,4-dicarboxamide**

**Example 55**

Reagent - 0.0295mL (0.253mmol) of morpholine carbonyl chloride  
4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(morpholin-4-  
ylcarbonyl)piperidine-4-carboxamide

**Example 56**

Reagent - 0.043g (0.253mmol) of methylphenylcarbamoyl chloride  
4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-4-hydroxy-N-1-methyl-N-1-  
phenylpiperidine-1,4-dicarboxamide

**Example 57**

Octyl-4-([4-(but-2-ynyloxy)phenyl]sulfonyl)methyl)-4-  
[(hydroxyamino)carbonyl]piperidine-1-carboxylate

Step A: A solution of 0.097g (0.23mmol) of N-(tert-butoxy)-4-([4-(2-butynyloxy)-  
phenyl]sulfonyl)methyl)-4-[4-(2-butynyloxy)phenyl]sulfonyl)methyl-4-piperidine-  
carboxamide (0.097g, 0.23mmol), octyl chloroformate (0.0495 ml, 0.253 mmol) and  
diisopropylethylamine (0.08 ml, 0.46 mmol) in 2mL of CH<sub>2</sub>Cl<sub>2</sub> was shaken at room  
temperature for 18h and then concentrated in vacuo.

Step B: same as StepB of Example 46.

The following hydroxamic acids were synthesized according to the procedures  
of Example 57 using the appropriate reagents.

**Example 58**

Reagent - 0.038mL (0.253mmol) of 4-methoxyphenyl chloroformate  
4-Methoxyphenyl-4-([4-(but-2-ynyloxy)phenyl]sulfonyl)methyl)-4-  
[(hydroxyamino) carbonyl]piperidine-1-carboxylate

-93-

**Example 59**

Reagent – 0.0323mL (0.253mmol) of benzenesulfonyl chloride  
4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(phenylsulfonyl)  
piperidine-4-carboxamide

5

**Example 60**

Reagent – 0.0457g (0.253mmol) of 1-methylimidazole-4-sulfonyl chloride  
4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-[(1-methyl-1H-  
imidazol-4-yl)sulfonyl]piperidine-4-carboxamide

10

**Example 61**

1-[2-(Benzylamino)acetyl]-4-([4-(but-2-ynyloxy)phenyl]sulfonyl)methyl)-N-  
hydroxypiperidine-4-carboxamide

Step A: A solution of N-(tert-butoxy)-4-([4-2-butynyloxy)phenyl] sulfonyl)methyl)-  
4-[4-(2-butynyloxy)phenyl]sulfonyl)methyl-4-piperidine carboxyamide (0.097g,  
15 0.23mmol), triethylamine (0.064mL, 0.64mmol), chloroacetyl chloride (0.064 ml, 0.64  
mmol), and 4-dimethylaminopyridine (0.002 g) in 2mL of CH<sub>2</sub>Cl<sub>2</sub> was shaken at room  
temperature for 18h. The solution was then treated with benzyl amine (0.075mL,  
0.69mmol) and was shaken for 18h and then concentrated in vacuo.

20

Step B: same as Step B of Example 46.

The following hydroxamic acids were synthesized according to the procedures  
of Example 61 using the appropriate amine reagents.

25

**Example 62**

Reagent – 0.060mL (0.69mmol) of morpholine  
4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(2-morpholin-4-  
ylacetyl)piperidine-4-carboxamide

30

**Example 63**

Reagent – 0.076mL (0.69mmol) of N-methylpiperazine

4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-[2-(4-methylpiperazin-1-yl)acetyl]piperidine-4-carboxamide

5

Example #	HPLC retention time (min.) <sup>1</sup>	MS <sup>2</sup> (M+H) <sup>+</sup>
46	1.85	395
47	2.20	498
48	1.71	458
49	2.11	472
52	2.30	438
53	2.85	486
57	3.80	523
58	2.98	517
54	2.87	466
55	2.33	480
56	2.84	500
59	2.92	507
60	2.40	511
50	2.67	471
51	2.64	477
61	2.14	514
62	1.86	494
63	1.84	507

<sup>1</sup> LC conditions: Hewlett Packard 1100; YMC ODS-A 4.6 mmx50 mm5 u column at 23°C; 10uL injection; Solvent A: 0.05% TFA/water; Solvent B: 0.05% TFA/acetonitrile; Gradient: Time 0: 98% A; 1 min: 98% A; 7 min: 10% A, 8 min: 98% A; Post time 1 min. Flow rate 2.5 mL/min; Detection: 220 and 254 nm DAD

10

<sup>2</sup>Mass Spec conditions: API-electrospray.

15

**Example 64**

1-Acetyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid  
hydroxamide

Step 1: 4-But-2-ynyloxybenzenesulfonyl fluoride:

To a solution of 4-but-2-ynyloxybenzenesulfonyl chloride (prepared from Example 30, step 4) (2.0 g, 8.18 mmol) in acetonitrile (10 ml) was added KF-CaF<sub>2</sub> (2.85g, 16.3 mmol) and the resulting mixture was stirred for 4 hours at room temperature. The reaction mixture was filtered and the filtrate was concentrated. The crude product was dissolved in EtOAc and washed with water. The organic layer

20



was dried over anhydrous  $\text{Na}_2\text{SO}_4$  and the solvent was removed to obtain 1.5 g (80%) of the product as solid.

Step 2: 4-(4-But-2-ynyloxybenzenesulfonyl)-piperidine-1,4-dicarboxylic acid tert-butyl ester methyl ester

To a solution of diisopropylamine (1.58 mL, 11.3 mmol) in THF (25 mL) at  $0^\circ\text{C}$  was added 2.5M n-BuLi (4.68 mL, 11.7 mmol) and the resulting mixture was stirred for 15 min at that temperature. The reaction mixture was cooled to  $-78^\circ\text{C}$  and a solution of 1-(tert-butyl)-4-methyl 1,4-piperidinecarboxylate (prepared from example 30, step 1) (2.67g, 11.0 mmol) in THF (40 mL) was added. The resulting mixture was stirred for 1h and a solution of 4-but-2-ynyloxy benzenesulfonyl fluoride (2.5g, 11.0 mmol) in THF (25 mL) was added into it. After stirring for 4h at rt, the reaction was quenched with satd. aqueous  $\text{NH}_4\text{Cl}$  solution and extracted with EtOAc, dried over anhydrous  $\text{Na}_2\text{SO}_4$ . The crude product was purified by silica gel chromatography to obtain 2.6g (53%) of the product as a solid;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  1.44(s, 9H), 1.87(m, 3H), 1.98(m, 2H), 2.32(m, 2H), 2.62(m, 2H), 3.74(s, 3H), 4.17(m, 2H), 4.74(m, 2H), 7.09(d, 2H,  $J=7.2$  Hz), 7.71(d, 2H,  $J=7.2$  Hz).

Step 3: 4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid methyl ester:

To a solution of product from step 2 (500 mg, 1.11 mmol) in methylene chloride (10 ml) was added 4M HCl (2 ml) and the resulting mixture was stirred for 2 hours at room temperature. The solid was filtered, washed with ether to obtain 410mg (95%) of the product as a solid.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.86(m, 3H), 2.52(m, 4H), 2.89(m, 2H), 3.52(m, 2H), 3.74(s, 3H), 4.74(m, 2H), 7.10(d, 2H,  $J=8.7$  Hz), 7.69(d, 2H,  $J=8.7$  Hz).

Step 4: 1-Acetyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid methyl ester

To a solution of product from step 3 (105 mg, 0.23 mmol) in methylene chloride (1 ml) was added triethylamine (93 mg, 0.92 mmol), acetyl chloride (18 mg,

0.23 mmol) followed by a catalytic amount of dimethylaminopyridine. The resulting mixture was stirred for 8 hours at room temperature, quenched with water and extracted with methylene chloride. The organic layer was dried over anhydrous sodium sulfate and concentrated to give 75 mg (80%) of the product as a solid.

5

Step 5: 1-Acetyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid:

A solution of the ester, from step 4 (240 mg, 0.61 mmol) and lithium hydroxide (18 mg, 0.75 mmol) in tetrahydrofuran/methanol/water (3:3:2) mixture was stirred at room temperature for 15 hours. The mixture was concentrated, acidified to pH 3-5 with 1N aqueous hydrochloric acid, and extracted with ethyl acetate. The organic layer was washed with brine and dried over anhydrous sodium sulfate. Removal of the solvent under vacuo gave the acid. Yield : 200 mg, (87%). <sup>1</sup>H NMR(300 MHz, acetone-d<sub>6</sub>):δ 1.84(t, 3H, J= 2.8 Hz), 1.90-2.05(m, 2H), 2.06(s, 3H), 2.25-2.51(m, 3H), 3.06(m, 1H), 4.04(m, 1H), 4.63(m, 1H), 4.86(q, 1H, J= 2.0).

15

Step 6: 1-Acetyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid hydroxamide:

To a solution of 1-acetyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid (180 mg, 0.48 mmol) in dimethylformamide was added hydroxybenzotriazol (77mg, 0.57 mmol) followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (127 mg, 0.66) and N-methylmorpholine (0.078 ml, 0.71 mmol). The resulting mixture was stirred for 1 h at room temperature when 50% aqueous hydroxylamine solution (0.145 ml, 2.37 mmol) was added and the mixture was stirred for 15 h at that temperature. The solvent was removed in *vacuo* and ethyl acetate/water was added to the crude product. The organic layer was separated and washed successively with 1N aqueous hydrochloric acid, water, saturated aqueous sodium bicarbonate, and water. The organic layer was dried over anhydrous sodium sulfate and the solvent was removed in *vacuo* to obtain 100 mg (53%) of the product as a solid. <sup>1</sup>H NMR(300 MHz, CDCl<sub>3</sub>):δ 1.64(m, 1H), 1.85(m, 3H), 1.99(s, 3H), 2.31(m, 4H), 2.83(m, 1H), 3.88(m, 1H), 4.41(m, 1H), 4.88(m, 2H), 7.16(d, 2H, J= 9.0 Hz), 7.66(d, 2H, J= 9.0 Hz), 9.20(m, 1H), 11.00(m, 1H); MS-ES: m/z395.2 (M+H)<sup>+</sup>.

30

**Example 65****1-Benzoyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid  
hydroxamide**

- 5 Step 1: 1-Benzoyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid  
methyl ester

To a solution of 4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic  
acid methyl ester, (400 mg, 1.03 mmol) in chloroform (10 ml) was added  
triethylamine (416 mg, 4.12 mmol), benzoyl chloride (144  $\mu$ l, 1.24 mmol) followed  
10 by a catalytic amount of dimethylaminopyridine. The resulting mixture was stirred  
for 15 hours at room temperature, quenched with water and extracted with methylene  
chloride. The organic layer was dried over anhydrous sodium sulfate and  
concentrated to give 375 mg (80%) of the product as a solid. MS-ES: m/z 456.1  
(M+H)<sup>+</sup>.

15

Step 2: 1-Benzoyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid

1-Benzoyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid  
was prepared, starting from 1-benzoyl-4-(4-but-2-ynyloxybenzenesulfonyl)-  
piperidine-4-carboxylic acid methyl ester (300 mg, 0.66 mmol) and lithium  
20 hydroxide (18 mg, 0.75 mmol). The resulting reaction mixture was worked up as  
outlined in Example 64, (step 5). Yield: 250 mg (86%) of the acid. HR - MS: m/z  
Calculated for C<sub>23</sub>H<sub>23</sub>NO<sub>6</sub>S 442.1319; Found 442.1317.

Step 3: 1-Benzoyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid  
25 hydroxamide

The general procedure for step 6 (Example 64) was followed using 1-benzoyl-  
4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid (100 mg, 0.23  
mmol) in dimethylformamide (2 ml), 1-hydroxybenzotriazole (36 mg, 0.27 mmol), 1-  
[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (62 mg, 0.32 mmol),  
30 N-methylmorpholine (0.038 ml, 0.35 mmol), and hydroxylamine (0.083 ml, 1.15  
mmol) to obtain 40 mg (38%) of the product as a solid. MS-ES: m/z 457.2 (M+H)<sup>+</sup>.

**Example 66****1-(4-Methoxybenzoyl)-4-(4-but-2-ynyloxy benzenesulfonyl)piperidine-4-carboxylic acid hydroxamide.**

Step 1: 1-(4-Methoxybenzoyl)-4-(4-but-2-ynyloxybenzenesulfonyl) piperidine-4-carboxylic acid methyl ester

To a solution of 4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid methyl ester (260 mg, 0.77 mmol) in chloroform (7 ml) was added triethylamine (311 mg, 3.08 mmol), 4-methoxybenzoyl chloride (158 mg, 0.92 mmol) followed by a catalytic amount of dimethylaminopyridine. The resulting mixture was stirred for 15 hours at room temperature, quenched with water and extracted with methylene chloride. The organic layer was dried over anhydrous sodium sulfate and concentrated to give 280 mg (75%) of the product as a solid. HR - MS: m/z Calculated for  $C_{25}H_{27}NO_7S$  486.1581; Found 486.1576.

Step 2: 1-(4-Methoxybenzoyl)-4-(4-but-2-ynyloxybenzenesulfonyl) piperidine-4-carboxylic acid.

1-(4-Methoxybenzoyl)-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid. was prepared following the procedure of Example 64 (step 5). Starting from 1-(4-methoxybenzoyl)-4-(4-but-2-ynyloxybenzenesulfonyl) piperidine-4-carboxylic acid methyl ester 250 mg, 0.52 mmol) in 4ml of tetrahydrofuran: methanol (1:1) and 1N sodium hydroxide (1.03 ml, 1.03 mmol) 150 mg of (62%) of the acid was isolated. HR - MS: m/z Calculated for  $C_{24}H_{25}NO_7S$  472.1425; Found 472.1426.

Step 3: 1-(4-Methoxybenzoyl)-4-(4-but-2-ynyloxybenzenesulfonyl) piperidine-4-carboxylic acid hydroxamide;

1-(4-Methoxybenzoyl)-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid hydroxamide was prepared following the procedure Example 64 (step 6). Starting from 1-(4-methoxybenzoyl)-4-(4-but-2-ynyloxybenzenesulfonyl)-piperidine-4-carboxylic acid (90 mg, 0.19 mmol) in dimethylformamide (2 ml), 1-hydroxybenzotriazole (31 mg, 0.23 mmol), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (51 mg, 0.27 mmol), N-methylmorpholine (0.031 ml,

-99-

0.28 mmol), and hydroxylamine (0.068 ml, 0.95 mmol), 70 mg (76%) of the product was isolated as solid. HR – MS: m/z Calculated for  $C_{24}H_{26}N_2O_7S$  487.1534; Found 487.1531.

5

**Example 67****4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(pyrrolidine-1-carbonyl)-4-piperidinecarboxamide**

Step1: 4-(4-but-2-ynyloxybenzenesulfonyl)-1-(pyrrolidine-1-carbonyl)-piperidine-4-carboxylic acid methyl ester

10 To a solution of 4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid methyl ester (400 mg, 1.03 mmol) in chloroform (10 ml) was added triethylamine (208 mg, 2.06 mmol), pyrrolidinecarbonyl chloride (206 mg, 1.54 mmol) followed by a catalytic amount of dimethylaminopyridine. The resulting mixture was stirred for 15 hours at room temperature, quenched with water and  
15 extracted with methylene chloride. The organic layer was dried over anhydrous sodium sulfate and concentrated to give 400 mg (87%) of the product as a solid; MS-ES: m/z 449.3 (M+H)<sup>+</sup>.

Step 2: 4-(4-but-2-ynyloxybenzenesulfonyl)-1-(pyrrolidine-1-carbonyl)-piperidine-4-carboxylic acid :

20 4-(4-But-2-ynyloxybenzenesulfonyl)-1-(pyrrolidine-1-carbonyl)-piperidine-4-carboxylic acid was prepared following the procedure of Example 64 (step 5). Starting from 4-(4-but-2-ynyloxybenzenesulfonyl)-1-(pyrrolidine-1-carbonyl)-piperidine-4-carboxylic acid methyl ester (250 mg, 0.52 mmol) in 4ml of  
25 tetrahydrofuran: methanol (1:1) and 1N sodium hydroxide (1.03 ml, 1.03 mmol), 150 mg of (62%) of the acid was isolated. HR – MS: m/z Calculated for  $C_{24}H_{25}NO_7S$  472.1425; Found 472.1426.

Step 3: 4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(pyrrolidine-1-carbonyl)-  
30 4-piperidinecarboxamide was prepared following the procedure Example 64 (step 6). Starting from 4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(pyrrolidine-1-carbonyl)-4-piperidinecarboxylic acid (255 mg, 0.23 mmol) in dimethylformamide (6

-100-

- ml), 1-hydroxybenzotriazole(96 mg, 0.71 mmol), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride(157 mg, 0.82 mmol), N-methylmorpholine (0.099 ml, 0.84 mmol), and hydroxylamine (0.181 ml, 2.8 mmol), 150 mg( 60%) of the product was isolated as a solid. HR – MS: m/z Calculated for  $C_{21}H_{27}N_3O_6S$  450.1693; Found 450.1692.

### Example 68

#### Ethyl 4-(4-but-2-ynyloxybenzenesulfonyl)-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxylate

- Step 1: 1-Ethyl 4-methyl 4-(4-but-2-ynyloxybenzenesulfonyl)-1,4-piperidine-dicarboxylate

To a solution of 4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid methyl ester (400 mg, 1.03 mmol) in chloroform (10 ml) was added sodium bicarbonate (865 mg, 10.3 mmol), ethylchloroformate(0.147 ml, 1.54 mmol). The resulting mixture was stirred for 15 hours at room temperature, quenched with water and extracted with methylene chloride. The organic layer was dried over anhydrous sodium sulfate and concentrated to give 425 mg (98%) of the product as a solid. MS-ES: m/z 424.4 (M+H)<sup>+</sup>.

- Step 2: 1-(Ethylcarbonyl)-4-(4-but-2-ynyloxybenzenesulfonyl)-1-piperidine-carboxylic acid

1-(Ethylcarbonyl)-4-(4-but-2-ynyloxybenzenesulfonyl)-1-piperidine-carboxylic acid was prepared following the procedure of Example 64 (step 5). Starting from 1-Ethyl 4-methyl 4-(4-but-2-ynyloxybenzenesulfonyl)-1,4-piperidine-dicarboxylate (400 mg, 0.95 mmol) in 8ml of tetrahydrofuran: methanol; water (1:1:0.5) and lithium hydroxide (50 mg, 2.04mmol), 340 mg of (88%) of the acid was isolated. HR – MS: m/z Calculated for  $C_{19}H_{23}NO_7S$  408.1122; Found 408.1126.

- Step 3: Ethyl 4-(4-but-2-ynyloxybenzenesulfonyl)-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxylate

Ethyl 4-(4-but-2-ynyloxybenzenesulfonyl)-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxylate was prepared following the procedure Example 64 (step 6).

-101-

Starting from 1-(Ethylcarbonyl)-4-(4-but-2-ynyloxybenzenesulfonyl)-1-piperidine-carboxylic acid (225 mg, 0.55 mmol) in dimethylformamide (6 ml), 1-hydroxy-benzotriazole (89 mg, 0.66 mmol), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (148 mg, 0.77 mmol), N-methylmorpholine (0.091 ml, 0.86 mmol), and  
5 hydroxylamine (0.168 ml, 2.75 mmol), 150 mg (64%) of the product was isolated as a solid. HR - MS: m/z Calculated for  $C_{19}H_{24}N_2O_7S$  425.1377; Found 425.1375.

### Example 69

#### 10 4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(trifluoromethyl)sulfonyl]-4-piperidinecarboxamide

Step 1: Methyl 4-(4-but-2-ynyloxybenzenesulfonyl)-1-[(trifluoromethyl)sulfonyl]-4-piperidinecarboxylate

To a solution of 4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid methyl ester (350 mg, 0.90 mmol) in chloroform (10 ml) was added triethyl-  
15 amine (182 mg, 1.81 mmol), trifluoromethanesulfonyl chloride (0.125 ml, 1.17 mmol) followed by a catalytic amount of dimethylaminopyridine. The resulting mixture was stirred for 15 hours at room temperature, quenched with water and extracted with methylene chloride. The organic layer was dried over anhydrous sodium sulfate and concentrated to give 245 mg (56%) of the product as a solid. HR - MS: m/z  
20 Calculated for  $C_{18}H_{20}F_3NO_7S_2$  484.0706; Found 484.0700.

Step 2: 4-(4-but-2-ynyloxybenzenesulfonyl)-1-[(trifluoromethyl)sulfonyl]-4-piperidinecarboxylic acid

4-(4-but-2-ynyloxybenzenesulfonyl)-1-[(trifluoromethyl)sulfonyl]-4-  
25 piperidinecarboxylic acid was prepared following the procedure of Example 64 (step 5). Starting from Methyl 4-(4-but-2-ynyloxybenzenesulfonyl)-1-[(trifluoromethyl)sulfonyl]-4-piperidinecarboxylate (225 mg, 0.47 mmol) in 8ml of tetrahydrofuran: methanol; water (1:1:0.5) and lithium hydroxide (24 mg, 0.98mmol), 175 mg of (80%) of the acid was isolated. MS-ES: m/z 468.1 (M-H).

30

-102-

Step 3: 4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(trifluoromethyl)-sulfonyl]-4-piperidinecarboxamide.

4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(trifluoromethyl)-sulfonyl]-4-piperidinecarboxamide was prepared following the procedure Example 64 (step 6). Starting from 4-(4-but-2-ynyloxybenzenesulfonyl)-1-[(trifluoromethyl)-sulfonyl]-4-piperidinecarboxylic acid (145 mg, 0.31 mmol) in dimethylformamide (3 ml), 1-hydroxybenzotriazole (50 mg, 0.37 mmol), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (83 mg, 0.47 mmol), N-methylmorpholine (0.051 ml, 0.47 mmol), and hydroxylamine (0.095 ml, 1.55 mmol), 90 mg (60%) of the product was isolated as a solid. HR - MS: m/z Calculated for  $C_{17}H_{19}F_3N_2O_7S_2$  485.0659; Found 485.0666.

#### Example 70

##### 4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(3-pyridinylcarbonyl)-4-piperidinecarboxamide

Step 1: Methyl 4-(4-but-2-ynyloxybenzenesulfonyl)-1-(3-pyridinylcarbonyl)-4-piperidinecarboxylate

To a solution of 4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid methyl ester (500 mg, 1.29 mmol) in methylene chloride (10 ml) was added triethylamine (443 mg, 4.39 mmol), nicotinyl chloride (276 ml, 1.55 mmol) followed by a catalytic amount of dimethylaminopyridine. The resulting mixture was stirred for 15 hours at room temperature, quenched with water and extracted with methylene chloride. The organic layer was dried over anhydrous sodium sulfate and concentrated to give 460 mg (78%) of the product as a solid. HR - MS: m/z Calculated for  $C_{23}H_{24}N_2O_6S$  457.1428; Found 457.1428.

Step 2 : 4-(4-But-2-ynyloxybenzenesulfonyl)-1-(3-pyridinylcarbonyl)-4-piperidine-carboxylic acid

4-(4-But-2-ynyloxybenzenesulfonyl)-1-(3-pyridinylcarbonyl)-4-piperidine carboxylic acid was prepared following the procedure of Example 64 (step 5). Starting from Methyl 4-(4-but-2-ynyloxybenzenesulfonyl)-1-(3-pyridinylcarbonyl)-



-103-

4-piperidinecarboxylate (430 mg, 0.94 mmol) in 8ml of tetrahydrofuran: methanol (1:1), and 1N sodium hydroxide (1.89 ml, 1.89 mmol) to obtain 235 mg(57%) of the acid. HR – MS: m/z Calculated for  $C_{22}H_{22}N_2O_6S$  443.1271; Found 443.1270.

- 5 Step 3: 4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(3-pyridinylcarbonyl)- 4-piperidinecarboxamide was prepared following the procedure Example 64 (step 6). Starting from 4-(4-But-2-ynyloxybenzenesulfonyl)-1-(3-pyridinylcarbonyl)-4-piperidinecarboxylic acid (195 mg, 0.44 mmol) in dimethylformamide (4 ml), 1-hydroxybenzotriazole (72 mg, 0.53 mmol), 1-[3-(dimethylamino)propyl]-3-ethyl-
- 10 carbodiimide hydrochloride(119 mg, 0.62 mmol), N-methylmorpholine (0.072 ml, 0.66 mmol), and hydroxylamine (0.135 ml, 2.2 mmol), 65 mg( 32%) of the product was isolated as a solid. HR – MS: m/z Calculated for  $C_{22}H_{23}N_3O_6S$  458.1380; Found 458.1373.

15

#### Example 71

#### 4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(2-thienylcarbonyl)- 4-piperidinecarboxamide

Step 1: Methyl 4-(4-but-2-ynyloxybenzenesulfonyl)-1-(2-thienylcarbonyl)- 4-piperidinecarboxylate

- 20 To a solution of 4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid methyl ester (500 mg, 1.29 mmol) in methylene chloride (10 ml) was added triethylamine (261 mg, 2.58 mmol), thiophenylcarbonyl chloride(227 mg, 1.55 mmol) followed by a catalytic amount of dimethylaminopyridine. The resulting mixture was stirred for 15 hours at room temperature, quenched with water and extracted with
- 25 methylene chloride. The organic layer was dried over anhydrous sodium sulfate and concentrated to give 480 mg (81%) of the product as a solid. HR – MS: m/z Calculated for  $C_{22}H_{23}NO_6S_2$  462.1040; Found 462.1039.

- Step 2: 4-(4-but-2-ynyloxybenzenesulfonyl)-1-(2-thienylcarbonyl)- 4-piperidine-
- 30 carboxylic acid

4-(4-but-2-ynyloxybenzenesulfonyl)-1-(2-thienylcarbonyl)-4-piperidine-carboxylic acid was prepared following the procedure of Example 64 (step 5).

-104-

Starting from Methyl 4-(4-but-2-ynyloxybenzenesulfonyl)-1-(2-thienylcarbonyl)-4-piperidinecarboxylate (435 mg, 0.94 mmol) in 8ml of tetrahydrofuran: methanol (1:1), and 1N sodium hydroxide (1.89 ml, 1.89 mmol) to obtain 360 mg(86%) of the acid. HR – MS: m/z Calculated for  $C_{21}H_{21}NO_6S_2$  448.0883; Found 448.0882.

5

Step 3: 4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(2-thienylcarbonyl)-4-piperidinecarboxamide

4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(2-thienylcarbonyl)-4-piperidinecarboxamide was prepared following the procedure Example 64 (step 6).

10 Starting from 4-(4-but-2-ynyloxybenzenesulfonyl)-1-(2-thienylcarbonyl)-4-piperidinecarboxylic acid (335 mg, 0.75 mmol) in dimethylformamide (7 ml), 1-hydroxybenzotriazole (121 mg, 0.90 mmol), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride(201 mg, 1.05 mmol), N-methylmorpholine (0.124 ml, 1.13 mmol), and hydroxylamine (0.229 ml, 3.75 mmol), 216 mg( 62%) of the  
15 product was isolated as a solid. HR – MS: m/z Calculated for  $C_{21}H_{22}N_2O_6S_2$  463.0992; Found 463.0988.

#### Example 72

20 4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(4-methoxyphenyl)sulfonyl]-4-piperidinecarboxamide.

Step 1 : Methyl 4-(4-but-2-ynyloxybenzenesulfonyl)-1-[(4-methoxyphenyl)sulfonyl]-4-piperidinecarboxylate

To a solution of 4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid methyl ester (500 mg, 1.29 mmol) in methylene chloride(10 ml) was added  
25 triethylamine (261 mg, 2.58 mmol), 4-methoxyphenylsulfonyl chloride(320 mg, 1.55 mmol) followed by a catalytic amount of dimethylaminopyridine. The resulting mixture was stirred for 15 hours at room temperature, quenched with water and extracted with methylene chloride. The organic layer was dried over anhydrous sodium sulfate and concentrated to give 590 mg (88%) of the product as a solid.

30 HR – MS: m/z Calculated for  $C_{24}H_{22}NO_8S_2$  522.1251; Found 522.1252.

-105-

Step 2: 4-(4-But-2-ynyloxybenzenesulfonyl)-1-[(4-methoxyphenyl) sulfonyl]-4-piperidinecarboxylic acid

4-(4-But-2-ynyloxybenzenesulfonyl)-1-[(4-methoxyphenyl)sulfonyl]-4-piperidine carboxylic acid was prepared following the procedure of Example 64 (step 5). Starting from methyl 4-(4-but-2-ynyloxybenzenesulfonyl)-1-[(4-methoxyphenyl)-sulfonyl]-4-piperidinecarboxylate (545 mg, 1.04 mmol) in 8ml of tetrahydrofuran: methanol (1:1), and 1N sodium hydroxide (2.09 ml, 2.09 mmol) to obtain 446 mg (85%) of the acid. HR – MS: m/z Calculated for  $C_{23}H_{25}NO_8S_2$  508.1094; Found 508.1073.

Step 3 : 4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(4-methoxyphenyl)-sulfonyl]-4-piperidinecarboxamide

4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(4-methoxyphenyl)-sulfonyl]-4-piperidinecarboxamide was prepared following the procedure Example 64 (step 6). Starting from 4-(4-But-2-ynyloxybenzenesulfonyl)-1-[(4-methoxyphenyl)-sulfonyl]-4-piperidinecarboxylic acid (402 mg, 0.79 mmol) in dimethylformamide (8 ml), 1-hydroxybenzotriazole (128 mg, 0.95 mmol), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (212 mg, 1.11 mmol), N-methylmorpholine (0.130 ml, 1.19 mmol), and hydroxylamine (0.242 ml, 3.95 mmol), 396 mg (96%) of the product was isolated as a solid. HR – MS: m/z Calculated for  $C_{23}H_{26}N_2O_8S_2$  523.1203; Found 523.1198.

### Example 73

4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxamide

Step 1: Methyl 4-(4-but-2-ynyloxybenzenesulfonyl)-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxylate

Methyl 4-(4-but-2-ynyloxybenzenesulfonyl)-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxylate was prepared following the procedure Example 64 (step 6). Starting from 4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid methyl ester (500 mg, 1.29 mmol) in dimethylformamide (10 ml), (2,2,5-trimethyl-1,3-dioxan-5-yl)carboxylic acid (224 mg, 1.29 mmol), 1-hydroxy-

-106-

benzotriazole (209 mg, 1.56 mmol), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (346 mg, 1.81 mmol), and N-methylmorpholine (0.212 ml, 1.94 mmol), to obtain 385 mg (59%) of the product as a solid. HR - MS: m/z Calculated for  $C_{25}H_{33}NO_8S$  508.2000; Found 508.1998.

5

Step 2: 4-(4-but-2-ynyloxybenzenesulfonyl)-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxylic acid

4-(4-But-2-ynyloxybenzenesulfonyl)-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)-carbonyl]-4-piperidinecarboxylic acid was prepared following the procedure of Example 40 (step 5). Starting from Methyl 4-(4-but-2-ynyloxybenzenesulfonyl)-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxylate (335 mg, 0.66 mmol) in 4ml of tetrahydrofuran: methanol (1:1), and 1N sodium hydroxide (1.3 ml, 1.3 mmol) to obtain 315 mg (97%) of the acid. HR - MS: m/z Calculated for  $C_{24}H_{31}NO_8S$  494.1843; Found 494.1835.

15

Step 3: 4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxamide:

4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxamide was prepared following the procedure of Example 64 (step 6). Starting from 4-(4-but-2-ynyloxybenzenesulfonyl)-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxylic acid (280 mg, 0.57 mmol) in dimethylformamide (6 ml), 1-hydroxybenzotriazole (92 mg, 0.57 mmol), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (153 mg, 0.80 mmol), N-methylmorpholine (0.094 ml, 0.85 mmol), and hydroxylamine (0.174 ml, 2.85 mmol), 180 mg (62%) of the product was isolated as a solid. HR - MS: m/z Calculated for  $C_{24}H_{32}N_2O_8S$  531.1771; Found 531.1768.

25

#### Example 74

4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[3-hydroxy-2-(hydroxymethyl)-2-methylpropanoyl]-4-piperidinecarboxamide

30

To a solution of product from Example 73 (150 mg, 0.29 mmol) in tetrahydrofuran (2 ml) was added 1N aqueous hydrochloric acid (2 ml) and the

-107-

resulting mixture was stirred for 4 hours. The organic layer was washed with sodium bicarbonate, brine and dried over anhydrous sodium sulfate. Solvent was removed to obtain 40 mg (29%) of the product. HR - MS: m/z Calculated for  $C_{21}H_{28}N_2O_8S$  469.1639; Found 469.1637.

5

### Example 75

#### **Tert-butyl 4-[[4-(2-butynyloxy)phenyl]sulfonyl]-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxalate**

Step 1: 1-(tert-butoxycarbonyl)-4-[[4-(2-butynyloxy)phenyl]sulfonyl]-4-

10 piperidinecarboxylic acid

A solution of 4-(4-but-2ynyloxybenzenesulfonyl)-piperidine-1,4-dicarboxylic acid tert-butyl ester methyl ester (from example 64, step 2) (15g, 33.2 mmol) in water (100 mL), methanol (50 mL) and tetrahydrofuran (50 mL) was treated with lithium hydroxide hydrate (2.73g, 66.4 mmol) and heated at reflux for 8h. The reaction

15 mixture was concentrated in vacuo and filtered through celite. To the filtrate was added aqueous 1N hydrochloric acid. A thick gum was obtained which was dissolved in dichloromethane and washed with water. Concentration of the organic phase gave a foam (14.9 g). Trituration with diethyl ether gave 1-(tert-butoxycarbonyl)-4-[[4-(2-butynyloxy)phenyl]sulfonyl]-4-

20 Electropray MS m/z 482 (M-H)<sup>-</sup>

Step 2 : Tert-butyl 4-[[4-(2-butynyloxy)phenyl]sulfonyl]-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxalate

Dimethylformamide (3.53 mL, 46 mmol) was added to a solution of oxalyl chloride (22.9 mL of a 2.0M solution in dichloromethane) in dichloromethane (25

25 mL) at 0°C. After 15 min a solution of 1-(tert-butoxycarbonyl)-4-[[4-(2-butynyloxy)phenyl]sulfonyl]-4-piperidinecarboxylic acid (10g, 22.9 mmol) in dimethylformamide was added and the reaction mixture was allowed to warm to room temperature. After 1h the reaction mixture was added to a mixture of hydroxylamine hydrochloride (16 g, 229 mmol), triethylamine (48 mL, 344 mmol), water (123 mL) and tetrahydrofuran

30 (500 mL) that had been stirring at 0°C for 15 min. The reaction was allowed to warm to room temperature. After 18h it was then diluted with ethyl acetate and washed with

-108-

saturated aqueous sodium bicarbonate (3X), then dried over potassium carbonate and concentrated in vacuo. Trituration with diethyl ether gave tert-butyl 4-{{[4-(2-butynyloxy)phenyl]sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxalate as a white powder (6.3g). <sup>1</sup>H NMR (dmso d<sub>6</sub>, 300 MHz) δ 1.38 (s, 9H, t-butyl), 1.6 – 1.7 (m, 2H, CHH), 1.85 (t, 3H, CH<sub>3</sub>, J = 2.2 Hz), 2.2 – 2.3 (m, 2H, CHH), 2.5 – 2.7 (m, 2H, NCHH), 3.9 – 4.0 (m, 2H, NCHH), 4.87 (q, 2H, OCH<sub>2</sub>, J = 2.2 Hz), 7.1 – 7.7 (m, 4H, ArH). Electrospray MS m/z 453 (M+H)<sup>+</sup>

#### Example 76

#### 4-{{[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxy-4-piperidinecarboxamide hydrochloride

To tert-butyl 4-{{[4-(2-butynyloxy)phenyl]sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxalate (prepared from Example 75) (6.3g, 13.9 mmol) was added 4N hydrochloric acid in dioxane. After 6h the reaction mixture was concentrated in vacuo. Methanol was added and the resulting mixture concentrated in vacuo. Dichloromethane was added and removed in vacuo (2X). Trituration with diethyl ether gave 4-{{[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxy-4-piperidinecarboxamide hydrochloride as a white powder (5.14g). <sup>1</sup>H NMR (dmso d<sub>6</sub>, 300 MHz) δ 1.86 (t, 3H, CH<sub>3</sub>, J = 2.2 Hz), 2.0 – 2.7 (m, 8H, CH<sub>2</sub>), 4.89 (q, 2H, OCH<sub>2</sub>, J = 2.2 Hz), 7.1 – 7.8 (m, 4H, ArH), 8.8 – 11.0 (m, 4H, NH<sub>2</sub>, NHOH). Electrospray MS m/z 353 (M+H)<sup>+</sup>

#### Example 77

#### Methyl ({4-{{[4-(2-butynyloxy)phenyl]sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinyl}methyl}benzoate hydrochloride

To 4-{{[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxy-4-piperidinecarboxamide hydrochloride (prepared from Example 76) (2.5 g, 6.43 mmol) and methyl 4-(bromomethyl)benzoate (1.62 g, 7.07 mmol) in methanol (100 mL) at 50°C was added triethylamine (2.25 mL, 16.1 mmol). After 30 min additional methanol (50 mL) was added. After 18 h the reaction mixture was concentrated in vacuo and 1N aqueous hydrochloric acid (10 mL) and water was added. The resulting solid was isolated and

-109-

to it was added methanol (20 mL) and 1N hydrochloric acid in diethylether (15 mL). To the resulting solution was added diethyl ether. Trituration of the precipitate gave methyl ({4-{{4-(2-butynyloxy)phenyl}sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinyl}methyl)benzoate hydrochloride as a white powder (2.4 g). <sup>1</sup>H NMR (dmso d<sub>6</sub>, 300 MHz) δ 1.85 (t, 3H, CH<sub>3</sub>, J = 2.2 Hz), 2.1 – 3.5 (m, 8H, CH<sub>2</sub>), 3.87 (s, 3H, OCH<sub>3</sub>), 4.40 (bd s, 2H, NCH<sub>2</sub>Ar), 4.89 (q, 2H, OCH<sub>2</sub>, J = 2.2 Hz), 7.1 – 8.1 (m, 8H, ArH), 9.3 – 11.2 (m, 3H, NH, NHOH). Electrospray MS m/z 501.5 (M+H)<sup>+</sup>

**Example 78**

10      **4-({4-{{4-(2-butynyloxy)phenyl}sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinyl}methyl)benzoic acid hydrochloride**

To methyl ({4-{{4-(2-butynyloxy)phenyl}sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinyl}methyl)benzoate hydrochloride (Prepared from example 77) (0.072g, 0.134 mmol) in methanol (1 mL) was added 1N aqueous sodium hydroxide (0.5 mL). After 18 h 1N aqueous hydrochloric acid (0.5 mL) was added and the reaction mixture concentrated in vacuo. Water was added and the precipitate triturated to give 4-({4-{{4-(2-butynyloxy)phenyl}sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinyl}methyl)benzoic acid hydrochloride as an off-white solid (0.040 g). <sup>1</sup>H NMR (dmso d<sub>6</sub>, 300 MHz) δ 1.85 (t, 3H, CH<sub>3</sub>, J = 2.2 Hz), 2.1 – 3.5 (m, 8H, CH<sub>2</sub>), 4.37 (bd s, 2H, NCH<sub>2</sub>Ar), 4.89 (q, 2H, OCH<sub>2</sub>, J = 2.2 Hz), 7.0 – 8.1 (m, 8H, ArH), 9.3 – 11.2 (m, 3H, NH, NHOH), 13.1 (bd s, 1H, COOH). Electrospray MS m/z 487.1 (M+H)<sup>+</sup>

**Example 79**

25      **1-[4-(Aminocarbonyl)benzyl]-4-{{4-(2-butynyloxy)phenyl}sulfonyl}-N-hydroxy-4-piperidinecarboxamide hydrochloride**

To methyl ({4-{{4-(2-butynyloxy)phenyl}sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinyl}methyl)benzoate hydrochloride (from example 77) (0.20 g) in methanol (10 mL) was added concentrated aqueous ammonium hydroxide (4 mL). After several weeks the reaction mixture was concentrated in vacuo and chromatographed on silica gel (methanol/dichloromethane) to give a white powder which was dissolved in dichloromethane and methanol. 1N Hydrochloric acid in

-110-

diethylether was added followed by additional diethylether. Trituration gave 1-[4-(aminocarbonyl)benzyl]-4-[[4-(2-butynyloxy)phenyl]sulfonyl]-N-hydroxy-4-piperidinecarboxamide hydrochloride as a white powder (0.106 g). <sup>1</sup>H NMR (DMSO d<sub>6</sub>, 300 MHz) δ 1.85 (t, 3H, CH<sub>3</sub>, J = 2.2 Hz), 2.2 – 3.5 (m, 8H, CH<sub>2</sub>), 4.33 (bd s, 2H, NCH<sub>2</sub>Ar), 4.89 (q, 2H, OCH<sub>2</sub>, J = 2.2 Hz), 7.1 – 8.0 (m, 8H, ArH), 7.47 (s, 1H, CONH), 8.04 (s, 1H, CONH), 9.35 (bd s, 1H, NHOH), 10.44 (bd s, 1H, NHOH), 11.1 (s, 1H, NH). Electrospray MS m/z 486.3 (M+H)<sup>+</sup>

### Example 80

#### 10      **Tert-butyl 4-[[4-(but-2-ynyloxy)phenyl]sulfinyl]-4- [(hydroxyamino)carbonyl]piperidine-1-carboxalate**

To tert-butyl 4-[[4-(but-2-ynyloxy)phenyl]sulfinyl]-4-[(hydroxyamino)carbonyl]piperidine-1-carboxalate (0.30 g) (obtained from example 14) in methanol (10 mL) was added 30% aqueous hydrogen peroxide (3 mL). After 3 days water and  
15      dichloromethane were added and the organic phase washed with aqueous Na<sub>2</sub>SO<sub>3</sub>. Concentration of the organic phase gave material which was dissolved in methanol (8 mL) and treated with 30% aqueous hydrogen peroxide. After several days workup as above gave tert-butyl 4-[[4-(but-2-ynyloxy)phenyl]sulfinyl]-4-[(hydroxyamino)-  
20      carbonyl]piperidine-1-carboxalate as a colorless foam (0.26 g). <sup>1</sup>H NMR (DMSO d<sub>6</sub>, 300 MHz) δ 1.38 (s, 9H, t-butyl), 1.5 – 1.7 (m, 2H, CHH), 1.85 (t, 3H, CH<sub>3</sub>, J = 2.2 Hz), 2.1 – 2.2 (m, 2H, CHH), 2.5 – 2.7 (m, 2H, NCHH), 3.8 – 4.0 (m, 2H, NCHH), 4.81 (q, 2H, OCH<sub>2</sub>, J = 2.2 Hz), 7.1 – 7.4 (m, 4H, ArH), 9.1 (s, 1H, NHOH), 10.8 (s, 1H, NHOH). Electrospray MS m/z 437.2 (M+H)<sup>+</sup>

### 25      **Example 81**

#### **4-(4-(But-2-ynyloxy-benzenesulfinyl)-piperidine-4-carboxylic acid hydroxamide hydrochloride**

To tert-butyl 4-[[4-(but-2-ynyloxy)phenyl]sulfinyl]-4-[(hydroxyamino)-  
carbonyl]piperidine-1-carboxalate (prepared from example 80) (0.26 g) was added 4N  
30      hydrochloric acid in dioxane (4 mL). After 1 h the reaction mixture was concentrated in vacuo. Methanol was added and removed in vacuo. Dichloromethane was added and removed in vacuo 3X to give 4-(4-(But-2-ynyloxy-benzenesulfinyl)-piperidine-4-



-111-

carboxylic acid hydroxamide hydrochloride as a yellow solid (0.19 g). <sup>1</sup>H NMR (dmso d<sub>6</sub>, 300 MHz) δ 1.86 (t, 3H, CH<sub>3</sub>, J = 2.2 Hz), 1.7 – 2.8 (m, 8H, CH<sub>2</sub>), 4.82 (q, 2H, OCH<sub>2</sub>, J = 2.2 Hz), 7.1 – 7.5 (m, 4H, ArH), 8.4 – 11.0 (m, 4H, NH<sub>2</sub>, NHOH). Electrospray MS m/z 337.2 (M+H)<sup>+</sup>

5

### Example 82

#### 1-(4-Bromo-benzyl)-4-(4-But-2-ynyloxy-benzenesulfinyl)-piperidine-4-carboxylic acid hydroxamide hydrochloride

To a solution of 4-(4-(But-2-ynyloxy-benzenesulfinyl)-piperidine-4-carboxylic acid hydroxamide hydrochloride (prepared from example 81) (0.162 g, 0.434 mmol) and 4-bromobenzylbromide (0.120 g, 0.478 mmol) in methanol was added triethylamine (0.13 mL, 0.91 mmol). After 4 h the reaction mixture was concentrated in vacuo and chromatographed on silica gel (methanol/dichloromethane) to give an oily solid which was dissolved in dichloromethane. To the solution was added 1N hydrochloric acid in ether (1 mL). Concentration in vacuo gave 1-(4-Bromo-benzyl)-4-(4-But-2-ynyloxy-benzenesulfinyl)-piperidine-4-carboxylic acid hydroxamide hydrochloride as a tan solid (0.102 g). <sup>1</sup>H NMR (dmso d<sub>6</sub>, 300 MHz) δ 1.85 (t, 3H, CH<sub>3</sub>, J = 2.2 Hz), 1.9 – 3.5 (m, 8H, CH<sub>2</sub>), 3.87 (s, 3H, OCH<sub>3</sub>), 4.3 (bd s, 2H, NCH<sub>2</sub>Ar), 4.82 (q, 2H, OCH<sub>2</sub>, J = 2.2 Hz), 7.0 – 7.8 (m, 8H, ArH), 9.2 – 11.1 (m, 3H, NH, NHOH). Electrospray MS m/z 505.1/507.2 (M+H)<sup>+</sup>

20

### Pharmacology

Representative compounds of this invention were evaluated as inhibitors of the enzymes MMP-1, MMP-9, MMP-13 and TNF-α converting enzyme (TACE). The standard pharmacological test procedures used, and results obtained which establish this biological profile are shown below.

25

#### Test Procedures for Measuring MMP-1, MMP-9, and MMP-13 Inhibition

These standard pharmacological test procedures are based on the cleavage of a thiopeptide substrates such as Ac-Pro-Leu-Gly(2-mercapto-4-methyl-pentanoyl)-Leu-Gly-OEt by the matrix metalloproteinases MMP-1, MMP-13 (collagenases) or MMP-9 (gelatinase), which results in the release of a substrate product that reacts

30

-112-

colorimetrically with DTNB (5,5'-dithiobis(2-nitro-benzoic acid)). The enzyme activity is measured by the rate of the color increase. The thiopeptide substrate is made up fresh as a 20 mM stock in 100% DMSO and the DTNB is dissolved in 100% DMSO as a 100 mM stock and stored in the dark at room temperature. Both the  
5 substrate and DTNB are diluted together to 1 mM with substrate buffer (50 mM HEPES pH 7.5, 5 mM  $\text{CaCl}_2$ ) before use. The stock of enzyme is diluted with buffer (50 mM HEPES, pH 7.5, 5 mM  $\text{CaCl}_2$ , 0.02% Brij) to the desired final concentration. The buffer, enzyme, vehicle or inhibitor, and DTNB/substrate are added in this order to a 96 well plate (total reaction volume of 200  $\mu\text{l}$ ) and the  
10 increase in color is monitored spectrophotometrically for 5 minutes at 405 nm on a plate reader and the increase in color over time is plotted as a linear line.

Alternatively, a fluorescent peptide substrate is used. In this test procedure, the peptide substrate contains a fluorescent group and a quenching group. Upon cleavage of the substrate by an MMP, the fluorescence that is generated is quantitated  
15 on the fluorescence plate reader. The assay is run in HCBC assay buffer (50mM HEPES, pH 7.0, 5 mM  $\text{Ca}^{+2}$ , 0.02% Brij, 0.5% Cysteine), with human recombinant MMP-1, MMP-9, or MMP-13. The substrate is dissolved in methanol and stored frozen in 1 mM aliquots. For the assay, substrate and enzymes are diluted in HCBC buffer to the desired concentrations. Compounds are added to the 96 well plate  
20 containing enzyme and the reaction is started by the addition of substrate. The reaction is read (excitation 340 nm, emission 444 nm) for 10 min. and the increase in fluorescence over time is plotted as a linear line.

For either the thiopeptide or fluorescent peptide test procedures, the slope of the line is calculated and represents the reaction rate. The linearity of the reaction  
25 rate is confirmed ( $r^2 > 0.85$ ). The mean ( $\bar{x} \pm \text{sem}$ ) of the control rate is calculated and compared for statistical significance ( $p < 0.05$ ) with drug-treated rates using Dunnett's multiple comparison test. Dose-response relationships can be generated using multiple doses of drug and  $\text{IC}_{50}$  values with 95% CI are estimated using linear regression.

-113-

Test Procedure for Measuring TACE Inhibition

Using 96-well black microtiter plates, each well receives a solution composed of 10  $\mu$ L TACE (final concentration 1 $\mu$ g/mL), 70 $\mu$ L Tris buffer, pH 7.4 containing 10% glycerol (final concentration 10 mM), and 10  $\mu$ L of test compound solution in DMSO (final concentration 1 $\mu$ M, DMSO concentration <1%) and incubated for 10 minutes at room temperature. The reaction is initiated by addition of a fluorescent peptidyl substrate (final concentration 100  $\mu$ M) to each well and then shaking on a shaker for 5 sec.

The reaction is read (excitation 340 nm, emission 420 nm) for 10 min. and the increase in fluorescence over time is plotted as a linear line. The slope of the line is calculated and represents the reaction rate.

The linearity of the reaction rate is confirmed ( $r^2 > 0.85$ ). The mean ( $\bar{x} \pm \text{sem}$ ) of the control rate is calculated and compared for statistical significance ( $p < 0.05$ ) with drug-treated rates using Dunnett's multiple comparison test. Dose-response relationships can be generated using multiple doses of drug and IC<sub>50</sub> values with 95% CI are estimated using linear regression.

Human Monocytic THP-1 Cell Differentiation Assay For Soluble Proteins(THP-1 Soluble Protein Assay)

Mitogenic stimulation of THP-1 cells cause differentiation into macrophage like cells with concomitant secretion of tumor necrosis factor (TNF- $\alpha$ ) and TNF receptor (TNF-R p75/80 and TNF-R p55/60) and Interleukin-8 (IL-8), among other proteins. In addition, non-stimulated THP-1 cells shed both the p75/80 and the p55/60 receptors over time. The release of membrane bound TNF- $\alpha$  and possibly TNF-R p75/80 and TNF-R p55/60, but not IL-8, is mediated by an enzyme called TNF- $\alpha$  converting enzyme or TACE. This assay can be used to demonstrate either an inhibitory or a stimulatory compound effect on this TACE enzyme and any cytotoxic consequence of such a compound.

THP-1 cells (from ATCC) are a human monocytic cell line which were obtained from the peripheral blood of a one year old male with acute monocytic

-114-

leukemia. They can be grown in culture and differentiated into macrophage like cells by stimulation with mitogens.

For the assay, THP-1 cells are seeded from an ATCC stock which was previously grown and frozen back at  $5 \times 10^6$ /ml/vial. One vial is seeded into a T25-flask with 16 mls of RPMI-1640 with glutamax (Gibco) media containing 10 % fetal bovine serum, 100 units/ml penicillin, 100  $\mu$ g/ml streptomycin, and  $5 \times 10^{-5}$  M 2-mercapto-ethanol (THP-1 media). Each vial of cells are cultured for about two weeks prior to being used for an assay and then are used for only 4 to 6 weeks to screen compounds. Cells are subcultured on Mondays and Thursdays to a concentration of  $1 \times 10^5$ /ml.

To perform an assay, the THP-1 cells are co-incubated in a 24 well plate with 50 ml/well of a 24 mg/ml stock of Lipopolysaccharide (LPS) (Calbiochem Lot# B13189) at 37°C in 5% CO<sub>2</sub> at a concentration of  $1.091 \times 10^6$  cells/ml (1.1 ml/well) for a total of 24 hours. At the same time, 50 ml/well of drug, vehicle or THP-1 media is plated in appropriate wells to give a final volume of 1.2 ml/well. Standard and test compounds are dissolved in DMSO at a concentration of 36 mM and diluted from here to the appropriate concentrations in THP-1 media and added to the wells at the beginning of the incubation period to give final concentrations of 100 mM, 30 mM, 10 mM, 3 mM, 1 mM, 300 nM, and 100 nM. Cell exposure to DMSO was limited to 0.1 % final concentration. Positive control wells were included in the experiment which had mitogen added but no drug. Vehicle control wells were included as well, which were identical to the positive control wells, except that DMSO was added to give a final concentration of 0.083%. Negative control wells were included in the experiment which had vehicle but no mitogen or drug added to the cells. Compounds can be evaluated for their effect on basal (non-stimulated) shedding of the receptors by replacing the LPS with 50 ml/well of THP-1 media. Plates are placed into an incubator set at 5% CO<sub>2</sub> and at 37°C. After 4 hours of incubation, 300 ml/well of tissue culture supernatant (TCS) is removed for use in an TNF- $\alpha$  ELISA. Following 24 hours of incubation, 700 ml/well of TCS is removed and used for analysis in TNF-R p75/80, TNF-R p55/60 and IL-8 ELISAs.

-115-

In addition, at the 24 hours timepoint, and the cells for each treatment group are collected by resuspension in 500 µl/well of THP-1 media and transferred into a FACS tube. Two ml/tube of a 0.5 mg/ml stock of propidium iodide (PI) (Boehringer Mannheim cat. # 1348639) is added. The samples are run on a Becton Dickinson FaxCaliber FLOW cytometry machine and the amount of dye taken up by each cell is measured in the high red wavelength (FL3). Only cells with compromised membranes (dead or dying) can take up PI. The percent of live cells is calculated by the number of cells not stained with PI, divided by the total number of cells in the sample. The viability values calculated for the drug treated groups were compared to the viability value calculated for the vehicle treated mitogen stimulated group ("vehicle positive control") to determine the "percent change from control". This "percent change from control" value is an indicator of drug toxicity.

The quantity of soluble TNF-a, TNF-R p75/80 and TNF-R p55/60 and IL-8 in the TCS of the THP-1 cell cultures are obtained with commercially available ELISAs from R&D Systems, by extrapolation from a standard curve generated with kit standards. The number of cells that either take up or exclude PI are measured by the FLOW cytometry machine and visualized by histograms using commercially available Cytologic software for each treatment group including all controls.

Biological variability in the magnitude of the response of THP-1 cell cultures requires that experiments be compared on the basis of percent change from "vehicle positive control" for each drug concentration. Percent change in each soluble protein evaluated from the "vehicle positive control" was calculated for each compound concentration with the following formula:

$$\% \text{ Change} = \frac{\text{pg/ml (compound)} - \text{pg/ml (veh pos control)}}{\text{pg/ml (veh pos control)} - \text{pg/ml (veh neg control)}} \times 100$$

For the soluble protein (TNF-a, p75/80, p55/60, IL-8) studies under stimulated conditions, the mean pg/ml of duplicate wells were determined and the results expressed as percent change from "vehicle positive control". For the soluble protein (p75/80 and p55/60 receptors) studies under non-stimulated conditions, the

-116-

mean pg/ml of duplicate wells were determined and the results expressed as percent change from "vehicle positive control" utilizing the following formula:

$$\% \text{ Change} = \frac{\text{pg/ml (compound neg control)} - \text{pg/ml (veh neg control)}}{\text{pg/ml (veh neg control)}} \times 100$$

IC50 values for each compound are calculated by non-linear regression analysis using customized software utilizing the JUMP statistical package.

For the cell viability studies, the viabilities (PI exclusion) of pooled duplicate wells were determined and the results expressed as % change from "vehicle positive control". The viability values calculated for the compound treated groups were compared to the viability value calculated for the "vehicle positive control" to determine "percent change from control" as below. This value "percent change from control" is an indicator of drug toxicity.

$$\% \text{ Change} = \frac{\% \text{ live cells (compound)}}{\% \text{ live cells (veh pos control)}} - 1 \times 100$$

#### References:

- Bjornberg, F., Lantz, M., Olsson, I., and Gullberg, U. Mechanisms involved in the processing of the p55 and the p75 tumor necrosis factor (TNF) receptors to soluble receptor forms. *Lymphokine Cytokine Res.* 13:203-211, 1994.
- Gatanaga, T., Hwang, C., Gatanaga, M., Cappuccini, F., Yamamoto, R., and Granger, G. The regulation of TNF mRNA synthesis, membrane expression, and release by PMA- and LPS-stimulated human monocytic THP-1 cells in vitro. *Cellular Immun.* 138:1-10, 1991.
- Tsuchiya, S., Yamabe, M., Yamaguchi, Y., Kobayashi, Y., Konno, T., and Tada, K. Establishment and characterization of a human acute monocytic leukemia cell line (THP-1). *Int. J. Cancer.* 26:1711-176, 1980.

Results of the above in vitro matrix metalloproteinase inhibition, TACE inhibition and THP standard pharmacological test procedures are given in Table 1 below.

5 Table 1:

Example #	TACE IC <sub>50</sub> <sup>a</sup>	THP @3μM <sup>b</sup>	MMP1 IC <sub>50</sub> <sup>c</sup>	MMP9 IC <sub>50</sub> <sup>a</sup>	MMP13 IC <sub>50</sub> <sup>a</sup>
1	65	46%	3.3	385	155
2	82	68%	2.57	164	39.6
3	55	34%	9	280	90
4	90	25%	2.6	148	47.3
5	188	30%	.3	400	180
6	393	NT	32.9%	58.9%	60%
7	123	21%	2.5	225	59.4
8	195	21%	4.7	218	72
9	166	12%	2.1	96.2	35.2
10					
11	98	7%	0.143	5.8	3.1
12	41.8%	+58	15	1000	1500
13	882	+69%	10	2000	800
14	67%	NT	26%	21%	32%
15	38%	NT	24%	25%	24%
16	46%	NT	10	2056	1465
17	139	0	10	2296	946
18	11.4	45%	10	1276	98
19	74	4%	10	10000	1321
20	30.1	47%	2643 nM	568	121
21	509	6%	> 10	3504	858
22	48.4%	5%	> 10	1814	1076
23	86.2	62%	3206 nM	160	64.4
24	180	41%	5671 nM	2078	463
25	695	3%	> 10	2740	1177
26	136	63%	1994 nM	25.1	22.1
27	168	13%	> 10	1542	426
28	150	13%	106 nM	15.4	5.3
29	127	13%	91 nM	16	4.7
30	102	0	>10	5899	2911
31	314	8	>10	>10000	>10000
32	100	0	>10	>10000	2752
33	327	8	>10	~5000	~10000
34	33	68	~10	1393	102
35	57	14	>10	>10000	~10000
36	4.8	58	3.9	2828	380
37	18	NT	8.6	8575	1024
38	19	53	~10	1443	279
39	11	NT	3.77	4275	809
40	63	41	0.707	425	36
41	37	60	2.677	1121	254

-118-

Example #	TACE IC <sub>50</sub> <sup>a</sup>	THP @3 $\mu$ M <sup>b</sup>	MMP1 IC <sub>50</sub> <sup>c</sup>	MMP9 IC <sub>50</sub> <sup>a</sup>	MMP13 IC <sub>50</sub> <sup>a</sup>
42	12	78%	~10	>10000	1627
43	13	66	~10	>10000	1640
44	56	49	35%	50.6%	2381
45	25	48	3.8	3584	423
46	105	NT	NT	NT	55.6%
47	227	NT	NT	NT	275
48	66	NT	NT	NT	1035
49	32.6	NT	NT	NT	1727
50	18.3	NT	NT	NT	352
51	21.5	NT	NT	NT	403
52	41.8	NT	NT	NT	3710
53	20.8	NT	NT	NT	1165
54	32.2	NT	NT	NT	104
55	70.7	NT	NT	NT	600
56	31.1	NT	NT	NT	3.2
57	694	NT	NT	NT	458
58	21.1	NT	NT	NT	179
59	53.3	NT	NT	NT	11.2
60	38.4	NT	NT	NT	8.0
61	56.4	NT	NT	NT	575
62	64.6	NT	NT	NT	64.6
63	66.6	NT	NT	NT	2229
64	47	30	4.076	560	136
65	73	3	3.532	448	105
66	106	73	2.768	430	81
67	72	18	2.028	853	345
68	77	10	2.249	1333	503
69	115	14	3.999	1246	499
70	87	62	2.963	639	113
71	113	14	3.117	811	183
72	221	56	4.157	1211	369
73	NT	NT	NT	NT	NT
74	132	39	4.338	963	287
75	134	-4	2.588	1951	284
76	201	26	4.503	7886	4019
77	114	52	2.187	149	349
78	64.5	64	1.051	364	73.7
79	70	83	2.420	129	50.6
80	90	-7	186	122	40
81	277	25	1.877	1035	593
82	135	16	257	125	62

a = nM or % inhibition

b = % inhibition

c =  $\mu$ M or % inhibition, unless otherwise indicated



-119-

Based on the results obtained in the standard pharmacological test procedures described above, the compounds of this invention were shown to be inhibitors of the enzymes MMP-1, MMP-9, MMP-13 and TNF- $\alpha$  converting enzyme (TACE) and are therefore useful in the treatment of disorders such as arthritis, tumor metastasis, tissue  
5 ulceration, abnormal wound healing, periodontal disease, graft rejection, insulin resistance, bone disease and HIV infection.

The compounds of this invention are also useful in treating or inhibiting pathological changes mediated by matrix metalloproteinases such as atherosclerosis, atherosclerotic plaque formation, reduction of coronary thrombosis from  
10 atherosclerotic plaque rupture, restenosis, MMP-mediated osteopenias, inflammatory diseases of the central nervous system, skin aging, angiogenesis, tumor metastasis, tumor growth, osteoarthritis, rheumatoid arthritis, septic arthritis, corneal ulceration, proteinuria, aneurysmal aortic disease, degenerative cartilage loss following traumatic joint injury, demyelinating diseases of the nervous system, cirrhosis of the liver,  
15 glomerular disease of the kidney, premature rupture of fetal membranes, inflammatory bowel disease, age related macular degeneration, diabetic retinopathy, proliferative vitreoretinopathy, retinopathy of prematurity, ocular inflammation, keratoconus, Sjogren's syndrome, myopia, ocular tumors, ocular angiogenesis/-neovascularization and corneal graft rejection.

20 Compounds of this invention may be administered neat or with a pharmaceutical carrier to a patient in need thereof. The pharmaceutical carrier may be solid or liquid.

Applicable solid carriers can include one or more substances which may also act as flavoring agents, lubricants, solubilizers, suspending agents, fillers, glidants,  
25 compression aids, binders or tablet-disintegrating agents or an encapsulating material. In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of  
30 the active ingredient. Suitable solid carriers include, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl

-120-

cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone, low melting waxes and ion exchange resins.

Liquid carriers may be used in preparing solutions, suspensions, emulsions, syrups and elixirs. The active ingredient of this invention can be dissolved or  
5 suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fat. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, colors, viscosity regulators, stabilizers or osmo-regulators.  
10 Suitable examples of liquid carriers for oral and parenteral administration include water (particularly containing additives as above, e.g., cellulose derivatives, preferable sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g., glycols) and their derivatives, and oils (e.g., fractionated coconut oil and arachis oil). For parenteral administration the carrier can  
15 also be an oily ester such as ethyl oleate and isopropyl myristate. Sterile liquid carriers are used in sterile liquid form compositions for parenteral administration.

Liquid pharmaceutical compositions which are sterile solutions or suspensions can be utilized by, for example, intramuscular, intraperitoneal or subcutaneous injection. Sterile solutions can also be administered intravenously. Oral  
20 administration may be either liquid or solid composition form.

The compounds of this invention may be administered rectally in the form of a conventional suppository. For administration by intranasal or intrabronchial inhalation or insufflation, the compounds of this invention may be formulated into an aqueous or partially aqueous solution, which can then be utilized in the form of an  
25 aerosol. The compounds of this invention may also be administered transdermally through the use of a transdermal patch containing the active compound and a carrier that is inert to the active compound, is non-toxic to the skin, and allows delivery of the agent for systemic absorption into the blood stream via the skin. The carrier may take any number of forms such as creams and ointments, pastes, gels, and occlusive  
30 devices. The creams and ointments may be viscous liquid or semi-solid emulsions of either the oil in water or water in oil type. Pastes comprised of absorptive powders

-121-

dispersed in petroleum or hydrophilic petroleum containing the active ingredient may also be suitable. A variety of occlusive devices may be used to release the active ingredient into the blood stream such as a semipermeable membrane covering a reservoir containing the active ingredient with or without a carrier, or a matrix  
5 containing the active ingredient. Other occlusive devices are known in the literature.


The dosage to be used in the treatment of a specific patient suffering a MMP or TACE dependent condition must be subjectively determined by the attending physician. The variables involved include the severity of the dysfunction, and the size, age, and response pattern of the patient. Treatment will generally be initiated  
10 with small dosages less than the optimum dose of the compound. Thereafter the dosage is increased until the optimum effect under the circumstances is reached. Precise dosages for oral, parenteral, nasal, or intrabronchial administration will be determined by the administering physician based on experience with the individual subject treated and standard medical principles.

15 Preferably the pharmaceutical composition is in unit dosage form, e.g., as tablets or capsules. In such form, the composition is sub-divided in unit dose containing appropriate quantities of the active ingredient; the unit dosage form can be packaged compositions, for example packed powders, vials, ampoules, prefilled syringes or sachets containing liquids. The unit dosage form can be, for example, a  
20 capsule or tablet itself, or it can be the appropriate number of any such compositions in package form.

-122-

## CLAIMS

1. A compound of formula



I

5

wherein:

- $R_1$  is hydrogen, aryl, heteroaryl, alkyl of 1-8 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, or -C4-C8-cycloheteroalkyl;
- $R_2$  and  $R_3$  are each, independently, hydrogen, alkyl of 1-6 carbon atoms, -CN, or -CCH<sub>3</sub>;
- $R_7$  is hydrogen, aryl, aralkyl, heteroaryl, heteroaralkyl, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 1-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, -C(O)- $R_1$ , -SO<sub>2</sub>- $R_1$ , -C(O)-NHR<sub>1</sub>, -C(O)NR<sub>5</sub>R<sub>6</sub>, -C(O)R<sub>1</sub>NR<sub>5</sub>R<sub>6</sub>, -C(O)-OR<sub>1</sub>, -C(NH)-NH<sub>2</sub>.
- $R_5$  and  $R_6$  are each, independently, hydrogen, alkyl of 1-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, aryl, aralkyl, heteroaryl, heteroaralkyl or -C4-C8-cycloheteroalkyl;
- $R_8$ ,  $R_9$ ,  $R_{10}$ , and  $R_{11}$  are each, independently, hydrogen, aryl or heteroaryl, cycloalkyl of 3-6 carbon atoms, -C4-C8-cycloheteroalkyl, alkyl of 1-18 carbon atoms, alkenyl of 2-18 carbon atoms, alkynyl of 2-18 carbon atoms; with the proviso that one of the pairs  $R_8$  and  $R_9$ ,  $R_9$  and  $R_{10}$  or  $R_{10}$  and  $R_{11}$ , together with the carbon atom or atoms to which they are attached, form a cycloalkyl ring of 3-6 carbon atoms, or a -C4-C8-cycloheteroalkyl ring;
- $R_{12}$  is hydrogen, aryl or heteroaryl, cycloalkyl of 3-6 carbon atoms, -C4-C8-cycloheteroalkyl, or alkyl of 1-6 carbon atoms;

25

-123-

A is O, S, SO, SO<sub>2</sub>, NR<sub>7</sub>, or CH<sub>2</sub>;

X is O, S, SO, SO<sub>2</sub>, NR<sub>7</sub>, or CH<sub>2</sub>;

Y is aryl or heteroaryl, with the proviso that A and X are not bonded to adjacent atoms of Y; and

n is 0-2;

or a pharmaceutically acceptable salt thereof.

2. A compound of Claim 1 wherein Y is phenyl, pyridyl, thienyl, furanyl, imidazolyl or triazolyl or thiadiazolyl.

3. A compound of Claim 1 selected from the group consisting of:
- 1-(4-Bromo-benzyl)-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide;
  - 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-methoxy-benzyl)-piperidine-4-carboxylic acid hydroxyamide;
  - 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-chloro-benzyl)-piperidine-4-carboxylic acid hydroxyamide;
  - 1-Benzyl-4-(4-but-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxamide;
  - 1-(4-Bromo-benzyl)-4-(4-pent-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide;
  - 1-(4-Bromo-benzyl)-4-(4-oct-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide;
  - 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-fluoro-benzyl)-piperidine-4-carboxylic acid hydroxyamide;
  - 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-cyano-benzyl)-piperidine-4-carboxylic acid hydroxamide;
  - 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(4-methyl-benzyl)-piperidine-4-carboxylic acid hydroxamide;

-124-

- 4-(4-But-2-ynyloxy-benzenesulfonyl)-1-(3,4-dichloro-benzyl)-piperidine-4-carboxylic acid hydroxamide;
- 1-(4-Bromo-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide;
- 5 1-(4-Bromo-benzyl)-4-[4-(4-piperidin-4-yl-but-2-ynyloxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide;
- 1-(4-Bromo-benzyl)-4-[4-(4-morpholin-4-yl-but-2-ynyloxy)-benzenesulfonyl]-piperidine-4-carboxylic acid hydroxyamide;
- 10 4-(4-But-2-ynyloxy-phenylsulfanyl)-4-hydroxycarbamoyl-piperidine-1-carboxylic acid tert-butyl ester;
- 4-(4-But-2-ynyloxy-phenylsulfanyl)-piperidine-4-carboxylic acid hydroxyamide
- 1-(4-Bromo-benzyl)-4-(4-but-2-ynyloxy-phenylsulfanyl)-piperidine-4-carboxylic acid hydroxyamide;
- 15 4-(4-But-2-ynyloxy-phenylsulfanylmethyl)-tetrahydro-pyran-4-carboxylic acid hydroxyamide;
- 4-(4-But-2-ynyloxy-benzenesulfonylmethyl)-tetrahydro-pyran-4-carboxylic acid hydroxyamide;
- 4-(4-But-2-ynyloxy-benzenesulfinylmethyl)-tetrahydro-pyran-4-carboxylic acid hydroxyamide;
- 20 4-{[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxytetrahydro-2H-pyran-4-carboxamide;
- 1-benzyl-4-{[3-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxy-4-piperidine carboxamide;
- 25 4-{[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxy-1-isopropyl-4-piperidine carboxamide;
- 4-{[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxy-1-(3-pyridinylmethyl)-4-piperidine carboxamide;
- 30 3-{[4-(2-Butynyloxy)phenyl]sulfonyl}-1-ethyl-N-hydroxy-3-piperidine-carboxamide;

-125-

- 3-{{4-(2-butynyloxy)phenyl}sulfonyl}-1-(4-chlorobenzyl)-N-hydroxy-3-piperidinecarboxamide;
- 4-{{4-(2-Butynyloxy)phenyl}sulfonyl}-1-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-piperidine-4-carboxylic acid hydroxyamide;
- 5 4-{{4-(2-Butynyloxy)phenyl}sulfonyl}-1-(3-pentanyl)-piperidine-4-carboxylic acid hydroxyamide;
- 1-(4-Methoxy-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide;
- 10 1-(4-Chloro-benzyl)-4-(4-prop-2-ynyloxy-benzenesulfonyl)-piperidine-4-carboxylic acid hydroxyamide;
- tert-butyl-4-({[4-(2-butynyloxy)phenyl]sulfanyl}methyl)-4-[(hydroxyamino)-carbonyl]-1-piperidinecarboxylate;
- 4-({[4-(But-2-ynyloxy)phenyl]thio}methyl)-N-hydroxypiperidine-4-carboxamide;
- 15 tert-Butyl-4-({[4-(2-butynyloxy)phenyl]sulfinyl}methyl)-4-[(hydroxyamino)-carbonyl]-1-piperidinecarboxylate;
- 4-[[[4-(2-Butynyloxy)phenyl]sulfinyl]methyl]-N-hydroxy-4-piperidine-carboxamide;
- tert-Butyl-4-({[4-(but-2-ynyloxy)phenyl]sulfonyl}methyl)-4-[(hydroxyamino)-carbonyl]piperidine-1-carboxylate;
- 20 tert-butyl-4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-4-[(hydroxyamino)-carbonyl]-1-piperidinecarboxylate;
- 1-Acetyl-4-[[[4-(2-butynyloxy)phenyl]sulfonyl]methyl]-N-hydroxy-4-piperidinecarboxamide;
- 25 1-(2-Butynyl)-4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-4-piperidinecarboxamide hydrochloride;

- N-1-(tert-Butyl)-4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-N-4-hydroxy-1,4-[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-N-4-hydroxy-1,4-l[sulfonyl]-methyl)-N-4-hydroxy-1,4-piperidinedicarboxamide;
- 5 Methyl 4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-4-[(hydroxyamino)-carbonyl]-1-piperidinecarboxylate;
- Benzyl 4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-4-[(hydroxyamino)-carbonyl]-1-piperidinecarboxylate;
- 1-Benzyl-4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-4-butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-4-piperidinecarboxamide;
- 10 4-({[4-(2-Butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxamide;
- 4-({[4-(2-Butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-1-[3-hydroxy-2-(hydroxymethyl)-2-methylpropanoyl]-4-piperidinecarboxamide;
- 1-[Amino(imino)methyl]-4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-4-l]-4-({[4-(2-butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-4-oxy)phenyl]sulfonyl}methyl)-N-hydroxy-4-piperidinecarboxamide;
- 15 4-({[4-(2-Butynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-1-(4-hydroxy-2-butynyl)-henyl]sulfonyl}methyl)-N-hydroxy-1-(4-hydroxy-2-butynyl)-4-piperidinecarboxamide;
- 20 4-({[4-(But-2-ynyloxy)phenyl]sulfonyl}methyl)-1-ethyl-N-hydroxypiperidine-4-carboxamide trifluoroacetic acid salt;
- 2-chloro-5-(chloromethyl)thiophene-4-({[4-(But-2-ynyloxy)phenyl]-sulfonyl}-methyl)-1-[(5-chlorothien-2-yl)methyl]-N-hydroxypiperidine-4-carboxamide trifluoroacetic acid salt;
- 25 4-({[4-(But-2-ynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-1-(pyridin-4-ylmethyl)piperidine-4-carboxamide trifluoroacetic acid salt;
- 4-({[4-(But-2-ynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-1-(pyridin-3-ylcarbonyl)piperidine-4-carboxamide trifluoroacetic acid salt;
- 1-Benzoyl-4-({[4-(but-2-ynyloxy)phenyl]sulfonyl}methyl)-N-hydroxy-
- 30 piperidine-4-carboxamide;



- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(thien-2-ylcarbonyl) piperidine-4-carboxamide;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-1-ethyl-N-4-hydroxypiperidine-1,4-dicarboxamide;
- 5 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-4-hydroxy-N-1-phenylpiperidine-1,4-dicarboxamide;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-1,N-1-diethyl-N-4-hydroxypiperidine-1,4-dicarboxamide;
- 10 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(morpholin-4-ylcarbonyl)piperidine-4-carboxamide;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-4-hydroxy-N-1-methyl-N-1-phenylpiperidine-1,4-dicarboxamide;
- Octyl-4-([4-(but-2-ynyloxy)phenyl]sulfonyl)methyl)-4-[(hydroxyamino)-carbonyl] piperidine-1-carboxylate;
- 15 4-Methoxyphenyl-4-([4-(but-2-ynyloxy)phenyl]sulfonyl)methyl)-4-[(hydroxyamino) carbonyl]piperidine-1-carboxylate;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(phenylsulfonyl)piperidine-4-carboxamide;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-[(1-methyl-1H-imidazol-4-yl)sulfonyl]piperidine-4-carboxamide;
- 20 1-[2-(Benzylamino)acetyl]-4-([4-(but-2-ynyloxy)phenyl]-sulfonyl)methyl)-N-hydroxypiperidine-4-carboxamide;
- 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-(2-morpholin-4-ylacetyl)piperidine-4-carboxamide;
- 25 4-([4-(But-2-ynyloxy)phenyl]sulfonyl)methyl)-N-hydroxy-1-[2-(4-methylpiperazin-1-yl)acetyl]piperidine-4-carboxamide;
- 1-Acetyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid hydroxamide;
- 1-Benzoyl-4-(4-but-2-ynyloxybenzenesulfonyl)piperidine-4-carboxylic acid hydroxamide;
- 30

-128-

- 1-(4-Methoxybenzoyl)-4-(4-but-2-ynyloxy benzenesulfonyl)piperidine-4-carboxylic acid hydroxamide;
- 4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(pyrrolidine-1-carbonyl)-4-piperidinecarboxamide;
- 5 Ethyl 4-(4-but-2-ynyloxybenzenesulfonyl)-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxylate;
- 4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(trifluoromethyl)sulfonyl]-4-piperidinecarboxamide;
- 4-(4-But-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(3-pyridinylcarbonyl)-4-piperidinecarboxamide;
- 10 4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-(2-thienylcarbonyl)-4-piperidinecarboxamide;
- 4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(4-methoxyphenyl)sulfonyl]-4-piperidinecarboxamide;
- 15 4-(4-but-2-ynyloxybenzenesulfonyl)-N-hydroxy-1-[(2,2,5-trimethyl-1,3-dioxan-5-yl)carbonyl]-4-piperidinecarboxamide;
- Tert-butyl 4-{[4-(2-butynyloxy)phenyl]sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinecarboxalate;
- 4-{[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxy-4-piperidinecarboxamide
- 20 hydrochloride;
- Methyl ({4-{[4-(2-butynyloxy)phenyl]sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinyl}methyl)benzoate hydrochloride;
- 4-({4-{[4-(2-butynyloxy)phenyl]sulfonyl}-4-[(hydroxyamino)carbonyl]-1-piperidinyl}methyl)benzoic acid hydrochloride;
- 25 1-[4-(Aminocarbonyl)benzyl]-4-{[4-(2-butynyloxy)phenyl]sulfonyl}-N-hydroxy-4-piperidinecarboxamide hydrochloride;
- Tert-butyl 4-{[4-(but-2-ynyloxy)phenyl]sulfonyl}-4-[(hydroxyamino)-carbonyl]piperidine-1-carboxalate;
- 4-(4-(But-2-ynyloxy-benzenesulfinyl)-piperidine-4-carboxylic acid
- 30 hydroxamide hydrochloride; and

-129-

1-(4-Bromo-benzyl)-4-(4-But-2-ynyloxy-benzenesulfinyl)-piperidine-4-carboxylic acid hydroxamide hydrochloride;  
and pharmaceutical salts thereof.

5           4.     A method of inhibiting pathological changes mediated by TNF- $\alpha$  converting enzyme (TACE) in a mammal in need thereof which comprises administering to said mammal a therapeutically effective amount of a compound having the formula I as defined in claim 1; or a pharmaceutically acceptable salt thereof.

10

5.     The method of Claim 4 wherein the condition treated is rheumatoid arthritis, graft rejection, cachexia, inflammation, fever, insulin resistance, septic shock, congestive heart failure, inflammatory disease of the central nervous system, inflammatory bowel disease or HIV infection.


15

6.     A pharmaceutical composition comprising a compound having the formula I as defined in claim 1 or a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable carrier.

20

7.     A process for preparing a compound of formula I as claimed in claim 1, which comprises one of the following:

a)     reacting a compound of formula



25     wherein n, X, Y, A, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>8</sub>, R<sub>9</sub>, R<sub>10</sub>, and R<sub>11</sub> are as defined in claim 1 or a reactive derivative thereof, with a compound of formula




wherein R<sub>12</sub> is as defined in claim 1, to give a compound of formula I;

-130-

or


b) deprotecting a compound of formula:



wherein  $n$ ,  $X$ ,  $Y$ ,  $A$ ,  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$  and  $R_{12}$ , are defined in claim 1, and  $R_{30}$  is a suitable protecting group such as *t*-butyl, benzyl, and trialkylsilyl, to give a corresponding compound of formula I

or

c) cleaving a resin supported hydroxamate derivative containing the group



wherein  $n$ ,  $X$ ,  $Y$ ,  $A$ ,  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_8$ ,  $R_9$ ,  $R_{10}$ , and  $R_{11}$ , are defined in claim 1 to give a compound of formula I wherein  $R_{12}$  is hydrogen;

or

d) resolving a mixture (e.g. racemate) of optically active isomers of a compound of formula I to isolate one enantiomer or diastereomer substantially free of the other enantiomer or diastereomers;

or

e) acidifying a basic compound of formula I with a pharmaceutically acceptable acid to give a pharmaceutically acceptable salt;

or

f) converting a compound of formula I having a reactive substituent group or site to a compound of formula I having a different substituent group or site.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 00/01864

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 C07D211/66 A61K31/445 C07D309/08 A61K31/35 C07D401/06 C07D409/06 C07D401/12 C07D413/04 C07D407/06 A61P19/00 A61P19/02		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 37877 A (AMERICAN CYANAMID COMPANY) 3 September 1998 (1998-09-03) cited in the application claims 1-3,14-19	3-7
P,X	WO 99 42436 A (AMERICAN CYANAMID COMPANY) 26 August 1999 (1999-08-26) claims 1-3,37-42	3-7
A	WO 97 24117 A (RHONE-POULENC RORER PHARMACEUTICALS INC.) 10 July 1997 (1997-07-10) cited in the application the whole document	3-7
<input type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : <div style="display: flex; justify-content: space-between;"> <div>           *A* document defining the general state of the art which is not considered to be of particular relevance            *E* earlier document but published on or after the international filing date            *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)            *O* document referring to an oral disclosure, use, exhibition or other means            *P* document published prior to the international filing date but later than the priority date claimed         </div> <div>           *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention            *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone            *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.            *Z* document member of the same patent family         </div> </div>		
Date of the actual completion of the international search		Date of mailing of the international search report
7 June 2000		20.06.00
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl Fax: (+31-70) 340-3016		Authorized officer  Hartrampf, G

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 00/01864

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although claims 4 and 5 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound.
2. ☒ Claims Nos.: 1,2 and 4 to 7 (partially)  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 00/01864

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1,2 and 4 to 7 (partially)

Present claims 1 and 2 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In fact, the claims contain so many options, variables, possible permutations and unclear definitions (see radicals R5, R6 and R7 which do not appear in formula (1)) that a lack of clarity and/or conciseness arises to such an extent as to render a meaningful search of the claims impossible. Consequently, the search has been carried out for those parts of the application which do appear to be clear and/or concise, namely those compounds recited in the examples, i.e. those mentioned in claim 3.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/01864

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9837877 A	03-09-1998	AU 6168698 A	18-09-1998
		AU 6436898 A	18-09-1998
		BR 9807803 A	22-02-2000
		EP 0973512 A	26-01-2000
		EP 0970046 A	12-01-2000
		NO 994124 A	26-10-1999
		NO 994125 A	26-10-1999
		WO 9838163 A	03-09-1998
WO 9942436 A	26-08-1999	AU 9120198 A	06-09-1999
WO 9724117 A	10-07-1997	AU 1529897 A	28-07-1997
		EP 0871439 A	21-10-1998
		JP 2000503012 T	14-03-2000